



**36th ANNUAL
RESIDENT
SCHOLARSHIP DAY**



**UNIVERSITY OF
SASKATCHEWAN**

Department of
Family Medicine

ABSTRACT BOOK
May 29, 2026

Department of Family Medicine
College of Medicine | University of Saskatchewan

medicine.usask.ca/familymedicine

May 12, 2026

I am looking forward to celebrating your scholarship achievements at the end of the month. This will be our 36th Annual event.


Asking and answering questions about practice is incredibly important for our patients and colleagues. I hope through the day; you will learn new things and be inspired to ask more questions about how we practice. The diversity of your questions re-enforces the breadth of our discipline.

Skilled evaluation of information, knowledge of and participation in the research process, and implementation, evaluation and adaptation of new processes are critical components of our ability to provide the best possible care to patients and communities.

To our graduating second-year and third-year residents, congratulations and all the best in your future practice. I also want to say thank-you to our team in the Scholarship Division as well as the faculty who have served as supervisors for these projects.

All the best in your future endeavours and keep asking questions!

Sincerely,



Kathy Lawrence
Provincial Head
Family Medicine

Congratulations on arriving at this stage of your Family Medicine Training!

As a former participant in the Resident Scholarship Day in Saskatchewan, I am aware of the feelings and emotions associated with completing a resident scholarship project. I am grateful we are able to come together to celebrate your success. Please know that your contributions to Family Medicine scholarship are valued and greatly appreciated.

The skills of research, scholarship and critical appraisal are essential and indispensable to all careers in Family Medicine. Medical information expands daily, and the rate of increase can be exponential in times of crisis. Considering this, your investment in your project has exposed you to skills that are crucial to your growth as a Family Physician.

It is my hope that as you read this collection of abstracts you are inspired to ask questions and find the answers to them. Scholarship is not just a list of publications. It is lifelong learning, inquiry, and critical appraisal of information.

Please join me in thanking those people who have made this moment possible: the Scholarship & Innovation Division, Faculty Advisors, Adjudicators, Operations personnel, and Award Sponsors which are only a partial list of the many important contributors. Thank you to all who make this day happen.

I would like to take this opportunity to wish the graduating FMRs all the best in the future and in your chosen careers.

Sincerely,



Sheila Smith, MD, CCFP (EM), FCFP
Postgraduate Program Director
Department of Academic Family Medicine
University of Saskatchewan



May 25, 2026

Dear Enhanced Skills Residents,

I would like to extend my sincere appreciation for your participation in the Family Medicine Resident Scholarship Day.

Scholarship is an important component of being an effective physician. It reflects our ongoing commitment to asking questions, seeking answers, and continually improving the care we provide to our patients, as well as the healthcare systems in which we work.

The Enhanced Skills year is demanding, and finding time to engage in formal scholarship is not easy. Your efforts in doing so are commendable, and I hope you found the process both valuable and rewarding. I look forward to learning more about your projects.

Congratulations on reaching this milestone and on the culmination of your hard work. I wish you all the best in your future careers and hope you will continue to incorporate scholarship into your practice.

Sincerely,

Martin Heroux
Enhanced Skills Program Director

May 29, 2026

Colleagues

On this occasion, the 36th Annual Resident Scholarship Day, I want to take this opportunity to recognize Residents, Research Officers, Coaches/Supervisors, Faculty, Staff and members of the various research teams for:

- all the hard work that has gone into making this possible;
- your commitment and perseverance given some of the challenges; and,
- for the many contributions that you have brought to these learning endeavours.

Since its inception in 1990, we have gathered together once a year to: celebrate our successes; learn about the scholarly questions that have been systematically answered over the past two years; ask and answer questions that will enhance our knowledge and understanding; and, provide reflections and feedback (peer-review). The Annual Scholarship Day in the Department of Family Medicine has evolved and grown over the years providing us with the opportunity to celebrate our academic achievements and to plan for the future.

Over the past 36 years, we have come a long way, but we must continue to transform to meet the needs of the peoples we serve; as well as the Accreditation Standards set by the College of Family Physicians of Canada.

Ian McWhinney (1926-2012) who was known as “the father of family medicine” transformed family medicine worldwide from a little-known subject or area of practice into an academic discipline with postgraduate training. Mahatma Gandhi stated, “you must be the change you wish to see in the world.” Paulo Freire stated in his book *Pedagogy of Hope* that “we must dare to invent the future.” Thus, improving practice provides opportunities for learning, answering questions and inventing the future each and every day.

I would also like to recognize the support that we receive from: the Division of Scholarship & Innovation; Department of Family Medicine; the College of Medicine; the University of Saskatchewan; the Saskatchewan College of Family Physicians; and the College of Family Physicians of Canada.

I wish each of you much success and the very best as you move forward in your chosen profession.

Yours sincerely,



Vivian R Ramsden, RN, PhD, MCFP (Hon.), FCAHS
Distinguished Professor & Director, Division of Scholarship & Innovation
Department of Family Medicine

36th ANNUAL RESIDENT SCHOLARSHIP DAY

DEPARTMENT OF ACADEMIC FAMILY MEDICINE

Friday, May 29, 2026
Prairieland Park – Hall C
Saskatoon, SK

- 0850 - 0900 Opening Remarks & Introductions
Dr. Vivian R Ramsden, Director, Division of Scholarship & Innovation
Dr. Sheila Smith, Program Director, Family Medicine Program
- 0900 – 0910 Assessing the Impact of the City of Prince Albert’s Alcohol Bylaw on Emergency Department Visits
Daniel Antonenko, Camille Mutukistna, Natalie Tinney, Jeffery DC Irvine, Lauren Rusk, Ananna Arna, Ahmed Faress, Anna Dinsdale
- 0910 – 0920 Understanding Emergency Department Use for Non-Emergent MSK Complaints in Moose Jaw, SK
Justin Hall, Anna Rucelli Ignacio, Mark Young, Brandon Thorpe, Adam Clay
- 0920 – 0930 Improving the Management of Chronic Obstructive Pulmonary Disease at the Family Medicine Unit in Regina
Ali Alfayyadh, Andrea Vasquez Camargo, Rejina Kamrul, Adam Clay, Radhika Marwah
- 0930 – 0940 What Family Physicians Need to Know About Creatine Use in Adolescents and Young Adults: A Scoping Review
Jason Fyfe, Boris Hencic, Katrina Sawchuk
- 0940 – 0950 Assessing Adherence for Counselling Non-Insulin-Dependent Type 2 Diabetics Regarding “Sick-Day Rules”: A Chart Review of 300 Chronic Disease Management (CDM) Patients at the Family Medicine Unit in Regina, Saskatchewan
Manraj Aujla, Manav Bhalla, Sanchsith Rajalingam, Rejina Kamrul, Adam Clay
- 0950 - 1000 Appropriate primary care prescribing and patient compliance with compression therapy for venous insufficiency
Maham Tasneem, Warda Zanib, Adam Clay, Rejina Kamrul, Abdalla Butt, David Kopriva
- 1000 - 1010 Evaluating Physicians’ Perceptions of Procedural Skills Taught in Saskatchewan’s Family Medicine Surgical Skills Curriculum
Chaudhry Masud, Husam Muftah, Jack Yassa, Clara Rocha Michaels, Adam Clay, Andrea Vasquez Camargo
- 1010 - 1040 Break**

- 1040 - 1050 A Rapid Review of Principles, Practices, and Challenges in Canada, Australia, and New Zealand Regarding the Teaching of Indigenous Health in Medical Education
Hamza Dani, Aaron Prystupa, Vivian R Ramsden, Rhonda Bryce
- 1050 - 1100 A Review of Preventative Care in HIV Care
Morgan Schatz, Jelyssa Luc, Cole Picot, Katrina Sawchuk
- 1100 - 1110 Accessing Healthcare and Affirming Care: Perspectives of 2SLGBTQIA+ Individuals in North Battleford, Saskatchewan
Jessica Kainth, Sarah Nazim, Michael Barnett, Ginger Ruddy, Rhonda Bryce
- 1110 - 1120 Investigating parenthood considerations and subsequent challenges while training in a Family Medicine Residency in Saskatchewan
Anum Ali, Aliaa El Tobgy, Eriny Tawedrous, Rhonda Bryce
- 1120 - 1130 Influence of prenatal hand expression on postpartum breastfeeding
Dilanjani Pietersz, Onyinyechi Nwoke, Anneme Dunhin, Rhonda Bryce
- 1130 - 1140 Evaluation of a Prenatal and Postpartum Educational Resource at West Winds Primary Health Care Centre
Abeer Nfaileh, Tracey Field, Katrina Sawchuk
- 1140 - 1150 Increasing Rates of Cervical Cancer Screening at a Rural Family Medicine Clinic
Oseihoria Alex-Ohunyon, Sonya Kim, Emmett Harrison, Adam Clay, Brenda Andreas
- 1150 - 1200 Effect of Non-Sterile Gloves on Infection Rate in Family Medicine Office-Based Procedures
Ololade Akinlabi, Sarina Sadeghi, Andrea Vasquez Carmargo, Adam Clay, Clara Rocha Michaels, Asma Gargoum
- 1200 - 1300 Lunch/Networking**
- 1300 - 1310 Artificial Intelligence in Primary Care: A Scoping Review of Perceptions, Adoption, and Emerging Implications of Generative AI
Carina Hoi Ying Chan, Aathmika Kirubaharan, Jared Oberkirsch, Udoka Okpalauwaekwe
- 1310 - 1320 Understanding Missed Appointments at West Winds Primary Health Centre: Perspectives and Opportunities
Stephanie Walls, Maa Quartey, Jason Hosain, Katrina Sawchuk
- 1320 - 1330 Exploring the Relationship Between Clinical Scheduling Practices, Resident Wellness, and Patient Care in Saskatchewan's Family Medicine Training Programs
Samantha Tso, Hadir Ismail, Mahmood Beheshti, Udoka Okpalauwaekwe

- 1330 - 1340 Exploration of Patient Perspectives of the Use of Artificial Intelligence Scribe in a Family Medicine Clinic
Riley Schaaf, Gezila De Beer, Mya Ajayi, Danielle Frost, Andries Muller, Katrina Sawchuk
- 1340 - 1350 Penicillin Allergy De-labelling at the Family Medicine Unit: A Quality Improvement Project
Kylee Kosokowsky, Shaina Templeton, Radhika Marwah, Debbie Bunka, Adam Clay
- 1350 - 1400 Dance Movement Therapy's Role in Chronic Disease Management: A Systematic Review
Vladyslav Liamprekht, Ai-Lien Le, Vipul Parekh, Rhonda Bryce, Breanna Davis
- 1400 - 1410 Enhanced Skills Program Evaluation: Graduate Survey of Career Satisfaction, Support Systems, and Work-Life Integration
Dinesh Kumar, Maria Siddiqui, Martin Heroux, Adam Clay
- 1410 - 1420 Pyxis-Programmed Anaphylaxis Kit to Improve Testing of Suspected Intraoperative Anaphylaxis: A Quality Improvement Project (FPA)
Yechan Kim, Vicky Loessin, Carla Flogan, Richard Schaan, Darcie Earle
- 1420 - 1430 Investigating Hypocalcemia in Trauma Patients: Trends in Detection and Treatment Timelines at RUH (FPA)
Chiru Gade, Troy Simpson, Matt Johnson, Darcie Earle
- 1430 - 1440 Do Visual Reminders Improve Appropriate ASA Prescribing for Preeclampsia Prevention?: A Quality Improvement Study
Yuli (Alex) Chen, Calista Lytle, Audrey Roane, Breanna Davis, Rhonda Bryce, Vivian R Ramsden
- 1440 - 1510 Break**
- 1510 - 1600 Recognition & Reflections - Dr. Vivian R Ramsden
Reflections - Dr. Jack Westfall, Dr. Mark Milne, Dr. Marcella Ogenchuk
Overall Research Award - Presented by the Saskatchewan College of Family Physicians:
 Lisa Bagonluri, Executive Director and Dr. Darcie McGonigle, President
Departmental Awards - Dr. Kathy Lawrence, Dr. Marty Heroux
Closing Remarks

Adjudicators

Jack Westfall – MD, MPH

John M. (Jack) Westfall grew up in Yuma, Colorado. He completed his MD and MPH at the University of Kansas School of Medicine, an internship in hospital medicine in Wichita, Kansas, and his Family Medicine Residency at the University of Colorado Rose Family Medicine Program. After joining the faculty at the University of Colorado's Department of Family Medicine, Dr Westfall started the High Plains Research Network, a geographic community and practice-based research network in rural and frontier Colorado. He practiced family medicine in several rural communities including Limon, Ft Morgan, and his hometown of Yuma. The HPRN was founded as a participatory practice-based research network, engaging clinicians and staff in the research. In 2003, the Community Advisory Council of farmers, ranchers, school teachers, and students joined the HPRN to fully engage patients and community members in all HPRN efforts. Clinicians and patients have been fully engaged partners in all HPRN research: defining research questions, assisting with methods and recruitment, leading interpretation and dissemination efforts, presenting and co-authoring manuscripts. The HPRN has received funding from the Centers for Disease Control, National Institutes of Health, Agency for Healthcare Research and Quality, and numerous state and local foundations. After retiring from the University of Colorado's School of Medicine, Jack worked for several years as the Director for Whole Person Care at Santa Clara County Health and Hospitals in San Jose, California. He also served for several years as the Director of the Robert Graham Center for policy research in primary care and family medicine in Washington DC. Returning to Colorado, he continued to consult and collaborate on primary care practice-based research, community-based participatory research, integrated primary care, behavioral health, and the interface between primary care, public, and community health. Jack practiced family medicine in his hometown, Yuma, Colorado until this year, when he followed his spouse to Boise, Idaho where he joined a local community health center.

Mark Milne, MSc, PhD – University of Saskatchewan, Saskatoon, SK

Dr. Mark Milne is the Research Facilitator for six departments in the College of Medicine at the University of Saskatchewan including the Department of Family Medicine. His role as a Research Facilitator ranges from meeting with students and faculty to discuss research ideas all the way to helping find and apply for grants. He has a MSc in Chemistry from the University of Saskatchewan, and a PhD in Chemistry and Medical Imaging from the University of Western Ontario. His PhD and research focused on the development and use of novel contrast agents for medical imaging along with the development of therapeutics for cancer treatment. Mark has been an adjudicator for the Department's Scholarship Day for the past few years and is excited to continue in this role at the 36th Annual Resident Scholarship Day.

Marcella Ogenchuk, RN, PhD – University of Saskatchewan, Saskatoon, SK

Dr Marcella Ogenchuk is a tenured Associate Professor in the University of Saskatchewan (USask)'s College of Nursing. She completed her BSN and Master of Nursing at USask, and her PhD from the College of Education, Department of Education Administration (USask). Dr. Ogenchuk has over 25 years of nursing experience in a variety of settings including acute, community including rural and remote settings in Saskatchewan. The focus of her practice and research is community-led. The population that she focuses on is children and youth.

Dr. Ogenchuk has been instrumental in developing pathways of care for urban elementary students using a multidisciplinary approach in the area of substance use prevention and oral health practice. She is working with communities in developing a tool for early identification of alcohol use disorder and has contributed to resources for families living with SUD (substance use disorder).

Master of Ceremonies – Jeffery DC Irvine, BSc, MD, MPH, CCFP

Dr. Jeff Irvine is a rural family physician in La Ronge, Saskatchewan, and faculty with the University of Saskatchewan's Department of Family Medicine. His practice has included primary care, emergency medicine, and obstetrics. He serves as Faculty Development Site Lead and Resident Scholarship Site Lead for the La Ronge Residency Training Program. His academic interests include rural and northern health systems, substance use in primary care, and community-partnered research, with a focus on supporting resident scholarship and locally relevant quality improvement.

Acknowledgements

The Division of Scholarship and Innovation in the Department of Family Medicine, University of Saskatchewan gratefully acknowledges the Saskatchewan College of Family Physicians and the Department of Family Medicine, University of Saskatchewan for the Resident Scholarship Awards.

Research Support

Dr. Rhonda Bryce, Adam Clay, Nicole Jacobson, Dr. Udoka Okpalauwaekwe, Dr. Katrina Sawchuk, Brenda Andreas and the Site Scholarship Leads (Drs. Mike Barnett, Mahmood Beheshti, Breanna Davis, Emmett Harrison, Jeff Irvine, Mark Lees, Evan Mah, Charleen Salmon, Andrea Vasquez Camargo, & Amanda Waldner).

Administrative Support

Adriana Cashwell, Jaime Markowski, Jaclyn Randall & the Program Administrators (Jalene Jepson, Tracy Arnold, Taryn Ashbee, Heidi Brown, Robyn Claypool, Kristen Huebner, Bobbie McLaughlin, Christina Morrice, Tamara Morrison, Jackie Powell, Janice Skilliter and Georgie Blackwell).

Prairieland Park

Kim Ferguson, Staff and Technical Support.

Assessing the Impact of the City of Prince Albert's Alcohol Bylaw on Emergency Department Visits

Daniel Antonenko, FMR II; Camille Mutukistna, FMR II; Natalie Timney, FMR II
Jeffery DC Irvine, BSc, MD, MPH, CCFP; Lauren Rusk, BHSc, MPH
Ananna Arna, BAsC, MSc; Ahmed Faress, MD MSc, FRCPC; Anna Dinsdale, BA, MPH

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Alcohol use is a major contributor to preventable morbidity and health system burden in Saskatchewan, where it is associated with more than 34,000 emergency department (ED) visits and hospitalizations annually. In June 2024, Prince Albert, Saskatchewan implemented a municipal bylaw reducing retail alcohol sale hours, creating a natural experiment to evaluate whether reduced alcohol availability was associated with changes in alcohol-related ED presentations.

Question(s): Was implementation of a municipal alcohol sale-hour restriction associated with a change in alcohol-related ED presentations in Prince Albert?

Methods/Methodology: We conducted a quasi-experimental interrupted time series study of alcohol-related ED presentations to Victoria Hospital from June 1, 2023 to May 31, 2025, including 12 months before and after policy implementation. ED visit data were obtained from the National Ambulatory Care Reporting System (NACRS). Alcohol-related visits were identified using alcohol-specific ICD-10-CA codes and structured chart review of selected non-alcohol-coded presentations. This study was approved by the University of Saskatchewan Research Ethics Board (Bio 1565) with Saskatchewan Health Authority operational approval (OA-UofS-5165).

Results/Findings: A total of 2,594 alcohol-related ED presentations were identified, representing 6.8% of all ED visits. Mean weekly alcohol-related ED presentations decreased from 27.0 before policy implementation to 22.7 after implementation, a 15.9% reduction. Interrupted time series analysis demonstrated a statistically significant immediate reduction in weekly alcohol-related ED presentations following policy implementation (level change coefficient -0.520 , standard error 0.167 ; $p=0.0018$).

Discussion: Implementation of a municipal alcohol sale-hour restriction in Prince Albert was associated with an immediate reduction in alcohol-related ED presentations. Alcohol involvement was also frequently identified in presentations not coded primarily as alcohol-related, suggesting administrative coding alone likely underestimates alcohol-related acute care burden.

Conclusions: Reduced municipal alcohol sale hours were associated with fewer alcohol-related ED presentations at Prince Albert's regional hospital.

Recommendations: Municipal regulation of alcohol retail availability should be considered a local policy tool to reduce alcohol-related harm and health system burden. Similar evaluations should be undertaken in other Canadian municipalities.

References:

1. Babor TF, Casswell S, Graham K, Huckle T, Livingston M, Rehm J, et al. Alcohol: No Ordinary Commodity-A Summary of the Third Edition. *Addiction*. 2022 Dec;117(12):3024-3036.
2. Bernal JL, Cummins S, Gasparrini A. Interrupted Time Series Regression for the Evaluation of Public Health Interventions: A Tutorial. *Int J Epidemiol*. 2017 Feb;46(1):348-355.
3. Canadian Institute for Health Information. Indicator Library: General Methodology Notes—Clinical Indicators [Internet]. Ottawa (ON): Canadian Institute for Health Information; 2023 Nov [cited 2026 May 14]. 27 p. Available from: <https://www.cihi.ca/sites/default/files/document/indicator-library-clinical-meth-notes-en.pdf>.
4. Canadian Institute for Health Information. Canadian Coding Standards for Version 2022 ICD-10-CA and CCI [Internet]. Ottawa (ON): Canadian Institute for Health Information; 2022 [cited 2026 May 14]. 771 p. Available from: https://secure.cihi.ca/free_products/canadian-coding-standards-2022-en.pdf.
5. Canadian Institute for Substance Use Research and Canadian Centre on Substance Use and Addiction. Canadian Substance Use Costs and Harms (2007-2014) [Internet]. Ottawa (ON): Canadian Centre on Substance Use and Addiction; 2018 [cited 2026 May 14]. 54 p. Available from: <https://www.ccsa.ca/sites/default/files/2019-04/CSUCH-Canadian-Substance-Use-Costs-Harms-Report-2018-en.pdf>.
6. Canadian Centre on Substance Use and Addiction. Canada’s Guidance on Alcohol and Health: Final Report [Internet]. Ottawa (ON): Canadian Centre on Substance Use and Addiction; 2023 [cited 2026 May 14]. 89 p. Available from: https://www.ccsa.ca/sites/default/files/2023-01/CCSA_Canadas_Guidance_on_Alcohol_and_Health_Final_Report_en.pdf.
7. City of Prince Albert. City of Prince Albert: Bylaw 13 of 2024 [Internet]. Prince Albert (SK): City of Prince Albert; 2024 Apr 22 [cited 2026 May 14]. 3 p. Available from: <https://www.citypa.ca/media/wggha21y/24-04-22-the-liquor-store-hours-bylaw-by1-13-2024.pdf>.
8. Hahn RA, Kuzara JL, Elder R, Brewer R, Chattopadhyay S, Fielding J, et al. Effectiveness of policies restricting hours of alcohol sales in preventing excessive alcohol consumption and related harms. *Am J Prev Med*. 2010 Dec;39(6):590-604.
9. King S, Paradis C, Reynolds J. An Evaluation of Psychoactive Substances That Bring Youth to the Emergency Department [Internet]. Ottawa (ON): Canadian Centre on Substance Use and Addiction; 2022 [cited 2026 May 14]. 64 p. Available from: https://www.ccsa.ca/sites/default/files/2022-12/Psychoactive%20Substances%20that%20Bring%20Youth%20to%20the%20Emergency%20Department_1.pdf.

10. Manthey J, Jacobsen B, Klinger S, Schulte B, Rehm J. Restricting Alcohol Marketing to Reduce Alcohol Consumption: A Systematic Review of the Empirical Evidence for One of the 'Best Buys'. *Addiction*. 2024 May;119(5):799-811.
11. McMillan GP, Hanson TE, Lapham SC. Geographic Variability in Alcohol-Related Crashes in Response to Legalized Sunday Packaged Alcohol Sales in New Mexico. *Accid Anal Prev*. 2007 Mar;39(2):252-257.
12. Liem Strategic Integration Inc. Alcohol Policy Review: Opportunities for Ontario Municipalities [Internet]. Sarnia (ON): Ontario Public Health Association; 2018 [cited 2026 May 14]. 68 p. Available from: <https://opha.on.ca/wp-content/uploads/2021/06/Alcohol-Policy-Review-Full-Report-Final-corrected.pdf>.
13. Popova S, Giesbrecht N, Bekmuradov D, Patra J. Hours and Days of Sale and Density of Alcohol Outlets: Impacts on Alcohol Consumption and Damage: A Systematic Review. *Alcohol Alcohol*. 2009 Sep-Oct;44(5):500-16.
14. Rehm J, Gmel GE Sr, Gmel G, Hasan OSM, Imtiaz S, Popova S, et al. The Relationship Between Different Dimensions of Alcohol Use and the Burden of Disease-An Update. *Addiction*. 2017 Jun;112(6):968-1001.
15. Statistics Canada [Internet]. Ottawa (ON): Government of Canada; c2026. Profile table, Census Profile, 2021 Census of Population – Prince Albert, City (CY), Saskatchewan; 2023 Nov 15 [cited 2026 May 14];[about 68 screens]. Available from: <https://www12.statcan.gc.ca/census-recensement/2021/dp-pd/prof/details/page.cfm?Lang=E&SearchText=prince%20albert&DGUIDlist=2021A00054715066&GENDERlist=1,2,3&STATISTIClist=1,4&HEADERlist=0>.
16. Treno AJ, Johnson FW, Remer LG, Gruenewald PJ. The Impact of Outlet Densities on Alcohol-Related Crashes: A Spatial Panel Approach. *Accid Anal Prev*. 2007 Sep;39(5):894-901.
17. Wagner AK, Soumerai SB, Zhang F, Ross-Degnan D. Segmented Regression Analysis of Interrupted Time Series Studies in Medication Use Research. *J Clin Pharm Ther*. 2002 Aug;27(4):299-309.
18. World Health Organization. Technical briefing for Appendix 3 of the Global Action Plan for Non-Communicable Diseases: Interventions to reduce the harmful use of alcohol [Internet]. Geneva (CE): World Health Organization; 2022 Nov 28 [cited 2026 May 15]. 5 p. Available from: <https://cdn.who.int/media/docs/default-source/ncds/mnd/technical-brief-alcohol.pdf>.

Understanding Emergency Department Use for Non-Emergent MSK Complaints in Moose Jaw, SK

Justin Hall, FMR II; Anna Ignacio, FMR II; Mark Young, FMR II
Brandon Thorpe, MB.BCh, CCFP, FLEX; Adam Clay, MSc

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Emergency departments (EDs) across Canada experienced overcrowding and prolonged wait times. In some regions, up to 25% of ED visits are for non-urgent musculoskeletal (MSK) presentations, which include sprains, back pain, and tendinopathies. However, no local data exists for Moose Jaw.

Question(s): What percentage of visits to the ED at Dr. F.H. Wigmore Regional Hospital between January 2022 and 2025 were for non-emergent MSK injuries within the scope of a physiotherapist?

Methods/Methodology: Data entered by the Dr. F.H. Wigmore Regional Hospital into the National Ambulatory Care Reporting Systems was used to identify visits between January 1, 2022, and January 31, 2025. Low acuity visits (CTAS 4 or 5) with an ICD diagnostic code within the scope of physiotherapy were included. This study received ethics approval from the University of Saskatchewan's Biomedical Research Ethics Board (Bio-5759) and Operational Approval from the Saskatchewan Health Authority (OA-UofS-5759 Exempt).

Results/Findings: 1,118 patients presented with diagnoses within the scope of physiotherapy practice, representing 1.6% of 71,881 total ED visits. The median age was 47 years (interquartile range: 34–66). Low back pain was the most frequent diagnosis (40.1%), and most patients (87.2%) were discharged home.

Discussion: Non-emergent MSK presentations accounted for a small proportion of ED visits, with low back pain being the most frequent diagnosis. This proportion is lower than that reported in prior literature, which has estimated a substantially larger share of ED utilization attributed to MSK conditions. This difference is likely due to methodological variation, as only CTAS 4–5 presentations were included and higher-acuity MSK cases were excluded. Although infrequent, these visits contribute to emergency department resource use, including physician assessment and diagnostic investigations.

Conclusions: Less than 2% of ED visits in Moose Jaw were for non-emergent MSK injuries within the physiotherapy scope. Although low in prevalence, these presentations represent a measurable component of ED utilization and contribute to overall service demand.

Recommendations: Strategies could include integrating allied healthcare professionals, such as physiotherapists, into ED workflows and developing alternative care pathways for non-urgent

MSK presentations. Future research could include all CTAS scores and evaluate healthcare costs of MSK visits.

References:

1. MacKay J, Atkinson P, Palmer E, Fraser J, Vaillancourt E, Howlett M, et al. Alternate Access to Care: A Cross Sectional Survey of Low Acuity Emergency Department Patients. *Cureus* [Internet]. 2017 Jun 23 [cited 20 Apr 2026];9(6):[12 p.]. Available from: <https://doi.org/10.7759/cureus.1385>.
2. Haas R, Gorelik A, Busija L, O'Connor D, Pearce C, Mazza D, Buchbinder R. Prevalence and Characteristics of Musculoskeletal Complaints in Primary Care: An Analysis from the Population Level and Analysis Reporting (POLAR) Database. *BMC Prim Care* [Internet]. 2023 Feb 4 [cited 20 Apr 2026];24(1):[10 p.]. Available from: <https://doi.org/10.1186/s12875-023-01976-z>.
3. Wiitavaara B, Fahlstrom M, Djupsjobacka M. Prevalence, Diagnostics and Management of Musculoskeletal Disorders in Primary Health Care in Sweden – An Investigation of 2000 Randomly Selected Patient Records. *J Eval Clin Pract*. 2017 Apr;23(2):325-332.
4. Baker B, Kessler K, Kaiser B, Waller R, Ingle M, Brambilla S, et al. Non-Traumatic Musculoskeletal Pain in Western Australian Hospital Emergency Departments: A Clinical Audit of the Prevalence, Management Practices and Evidence-to-Practice Gaps. *Emerg Med Australas*. 2019 Dec;31(6):1037-1044.
5. Bird S, Thompson C, Williams KE. Primary Contact Physiotherapy Services Reduce Waiting and Treatment Times for Patients Presenting with Musculoskeletal Conditions in Australian Emergency Departments: An Observational Study. *J Physiother*. 2016 Oct;62(4):209-214.
6. Lebec MT, Cernohous S, Tenbarga L, Gest C, Severson K, Howard S. Emergency Department Physical Therapist Service: A Pilot Study Examining Physician Perceptions. *Internet J Allied Health Sci Pract* [Internet]. 2010 Jan 1 [cited 2025 Feb 10];8(1):[12 p.]. Available from: <https://nsuworks.nova.edu/ijahsp/vol8/iss1/8/>.
7. Pugh A, Roper K, Magel J, Fritz J, Colon N, Robinson S, et al. Dedicated Emergency Department Physical Therapy Is Associated with Reduced Imaging, Opioid Administration, and Length of Stay: A Prospective Observational Study. *PLoS One* [Internet]. 2020 Apr 23 [cited 2025 Feb 10];15(4):[12 p.]. Available from: <https://doi.org/10.1371/journal.pone.0231476>.
8. Gurley KL, Blodgett MS, Burke R, Shapiro NI, Edlow JA, Grossman SA. The Utility of Emergency Department Physical Therapy and Case Management Consultation in Reducing Hospital Admissions. *J Am Coll Emerg Physicians Open*. 2020 Jun;1(5):880-886.
9. Sutton M, Govier A, Prince S, Morphett M. Primary-Contact Physiotherapists Manage a Minor Trauma Caseload in the Emergency Department Without Misdiagnoses or Adverse Events: An Observational Study. *J Physiother*. 2015 Apr;61(2):77-80.
10. Ronksley PE, Liu EY, McKay JA, Kobewka DM, Rothwell DM, Mulpuru S, Forster AJ. Variations in Resource Intensity and Cost Among High Users of the Emergency Department. *Acad Emerg Med*. 2016 Jun;23(6):722-730.
11. Ojha HA, Snyder RS, Davenport TE. Direct Access Compared with Referred Physical Therapy Episodes of Care: A Systematic Review. *Phys Ther*. 2014 Jan;94(1):14–30.
12. Hon S, Ritter R, Allen DD. Cost-Effectiveness and Outcomes of Direct Access to Physical Therapy for Musculoskeletal Disorders Compared to Physician-First Access in the United States: A Systematic Review and Meta-Analysis. *Phys Ther* [Internet]. 2021 Jan 4 [cited 2026 Apr 27];101(1):[11 p.]. Available from: <https://doi.org/10.1093/ptj/pzaa201>.

Improving the Management of Chronic Obstructive Pulmonary Disease (COPD) at the Family Medicine Unit in Regina

Ali Alfayyadh, FMR II; Andrea Vasquez Camargo, MD, CCFP, MSc;
Rejina Kamrul, MBBS, CCFP; Adam Clay, MSc;
Radhika Marwah, MBBS, CCFP, MSc-AMH (Addictions)

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Chronic obstructive pulmonary disease (COPD) affects 10% of Canadians aged ≥ 35 years¹. At the Family Medicine Unit (FMU) in Regina, Saskatchewan, the number of spirometry-confirmed COPD patients and adherence to the 2023 Canadian Thoracic Society (CTS) recommended baseline pharmacotherapy were unknown.

Question(s): This quality improvement project aims to determine the percentage of patients with documented COPD diagnosis at the FMU with confirmatory spirometry within the past five years and to assess the proportion receiving baseline long-acting bronchodilator (LABD) therapy per the 2023 CTS guidelines.

Methods/Methodology: Patients over the age of 50 were identified using the Canadian Primary Care Sentinel Surveillance Network case definitions for COPD, with the addition of asthma-COPD overlap diagnosis. A retrospective chart audit was conducted using MedAccess electronic medical records, reviewing spirometry status (2020–2024) and pharmacotherapy. The project received exemption from the University of Saskatchewan’s Biomedical Research Ethics Board (E-Bio-080) and Operational Approval from the Saskatchewan Health Authority (E-Bio-080 Exempt).

Results/Findings: Of the 270 patients identified, 135 were included in the chart review, of whom 132 were active. Spirometry confirmation was documented for 51 patients (38.6%). Among these, 16 had Chronic Disease Management (CDM) visits in 2024, of whom 14 (87.5%) were prescribed LABD therapy. Overall, 42 patients (82.4%) were on LABD therapy. Among the 81 non-confirmed cases, 48 had spirometry performed, with 27 showing no evidence of COPD, 20 having uninterpreted results, and 1 interpreted as a borderline for obstruction.

Discussion: This project highlighted a low spirometry confirmation rate of 38.6%. Adherence to LABD therapy was relatively good overall amongst confirmed patients (82.4%) and even higher among those with CDM visits within 2024 (87.5%). A notable finding was that 20 spirometry tests remained uninterpreted. These results reflect the challenges of diagnosing COPD in primary care.

Conclusions: The project established key baseline data for the FMU. While spirometry confirmation remains low and many uninterpreted, adherence to LABD therapy is relatively

strong especially among patients with recent CDM visits. These findings highlight opportunities for quality improvement interventions.

Recommendations: Future interventions may include pharmacotherapy adjustment recommendations to the most responsible physicians, enhanced spirometry interpretation pathways and documentation, and optimizing CDM template use.

References:

1. Public Health Agency of Canada [Internet]. Ottawa (ON): Government of Canada; c2026. Canadian Chronic Disease Surveillance System (CCDSS) Data Tool [Internet]; 2025 [cited 2025 Mar 13];[about 5 screens]. Available from: <https://health-infobase.canada.ca/ccdss/data-tool/index?G=00&V=7>.
2. Bailey KL. The Importance of the Assessment of Pulmonary Function in COPD. *Med Clin North Am.* 2012 Jul;96(4):745-752.
3. Bourbeau J, Bhutani M, Hernandez P, Aaron SD, Beaulac MF, Kermel SB, et al. 2023 Canadian Thoracic Society Guideline on Pharmacotherapy in Patients with Stable COPD. *Chest.* 2023 Nov;164(5):1159-1183.
4. Canadian Primary Care Sentinel Surveillance Network. CPCSSN Case Definitions 2024-Q4 [Internet]. Kingston (ON): Queen's University; 2019 May 1 [updated 2025 Jan 21; cited 2026 May 15]. 59 p. Available from: <https://cpcssn.ca/wp-content/uploads/2025/03/CPCSSN-Case-Definitions-2024-Q4.pdf>.
5. Diab N, Gershon AS, Sin DD, Tan WC, Bourbeau J, Boulet LP, Aaron SD. Underdiagnosis and Overdiagnosis of Chronic Obstructive Pulmonary Disease. *Am J Respir Crit Care Med.* 2018 Nov;198(9):1130-1139.
6. Gershon A, Mecredy G, Croxford R, To T, Stanbrook MB, Aaron SD. Outcomes of Patients with Chronic Obstructive Pulmonary Disease Diagnosed with or without Pulmonary Function Testing. *CMAJ.* 2017 Apr;189(14):E530-538.
7. Blazer AJ, Stanbrook MB. How Can Canada's Health Systems Improve Care for People with Chronic Obstructive Pulmonary Disease? *CMAJ.* 2023 Sept;195(35):E1182-1183.

What Family Physicians Need to Know About Creatine Use in Adolescents and Young Adults: A Scoping Review

Jason Fyfe, FMR II; Boris Hencic, MD, CCFP
Katrina Sawchuk, MEd, PhD

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Creatine is one of the most widely used ergogenic supplements among adolescents and young adults, particularly those involved in competitive sports. Despite its popularity, many family physicians report uncertainty regarding its safety, dosing, and benefits in this population, highlighting the need for a synthesis of current evidence to support primary care counselling.

Question(s): What evidence exists regarding the safety, dosing, and potential benefits of creatine use in individuals aged 15-25, and what key information should family physicians understand when advising patients?

Methods/Methodology: A Scoping Review was conducted using Covidence. Google Scholar, Embase, Ovid MEDLINE, and PubMed were searched for English-language randomized controlled trials and relevant reviews published within the last 20 years involving healthy, non-pregnant individuals aged 15-25. Due to the presence of all study materials being in the public domain, formal ethics approval was not required.

Results/Findings: Evidence suggests creatine supplementation is generally well tolerated in healthy adolescents and young adults when used at recommended doses. Common regimens include a loading phase of 0.3 g/kg/day for 5-7 days followed by maintenance dosing of 3-5 g/day, though lower-dose strategies without loading also appear effective. Most studies did not evaluate safety as a primary outcome, limiting the strength of safety conclusions. A consistent effect is weight gain related to increased intracellular water retention. Gastrointestinal complaints and muscle cramping have been reported, though these effects are not clearly attributable to creatine. Some studies suggest supplement use among adolescents, including creatine, may be associated with higher-risk of using other performance-enhancing substances such as anabolic steroids. Reported benefits include modest improvements in muscle strength, lean body mass, and high-intensity exercise performance. Limited emerging evidence also suggested a possible role for creatine in treating depression in adolescents.

Conclusions: Creatine use is common among adolescents and young adults, though discussions within family medicine appear limited. Current evidence suggests creatine is generally safe and well tolerated when used appropriately; however, long-term safety data remain limited.

Recommendations: Family physicians should routinely ask about supplement use, screen for concurrent higher-risk substances, and remain up to date with emerging literature to support informed counselling.

References:

1. Holland-Hall C. Performance-Enhancing Substances: Is Your Adolescent Patient Using? *Pediatr Clin North Am*. 2007 Aug;54(4):651–662.
2. Castillo EM, Comstock RD. Prevalence of Use of Performance-Enhancing Substances Among United States Adolescents. *Pediatr Clin North Am*. 2007 Aug;54(4):663–675.
3. Wiens K, Erdman KA, Stadnyk M, Parnell JA. Dietary Supplement Usage, Motivation, and Education in Young, Canadian Athletes. *Int J Sport Nutr Exerc Metab*. 2014 Dec;24(6):613–622.
4. Lattavo A, Kopperud A, Rogers PD. Creatine and Other Supplements. *Pediatr Clin North Am*. 2007 Aug;54(4):735–760.
5. Gradidge P, Coopoo Y, Constantinou D. Prevalence of Performance-Enhancing Substance Use by Johannesburg Male Adolescents Involved in Competitive High School Sports. *Arch Exerc Health Dis*. 2011 Jan;2(2):114–119.
6. Judge LW, Petersen JC, Craig BW, Hoover DL, Holtzclaw KA, Leitzelar BN, et al. Creatine Usage and Education of Track and Field Throwers at National Collegiate Athletic Association Division I Universities. *J Strength Cond Res*. 2015 Jul;29(7):2034–2040.
7. Kreider RB, Kalman DS, Antonio J, Ziegenfuss TN, Wildman R, Collins R, et al. International Society of Sports Nutrition Position Stand: Safety and Efficacy of Creatine Supplementation in Exercise, Sport, and Medicine. *J Int Soc Sports Nutr* [Internet]. 2017 Jun 13 [cited 2026 May 15];14(1):[18 p.]. Available from: <https://doi.org/10.1186/s12970-017-0173-z>.
8. Herriman M, Fletcher L, Tchaconas A, Adesman A, Milanaik R. Dietary Supplements and Young Teens: Misinformation and Access Provided by Retailers. *Pediatrics* [Internet]. 2017 Feb 1 [cited 2026 May 15];139(2):[9 p.]. Available from: <https://doi.org/10.1542/peds.2016-1257>.
9. Huerta Ojeda Á, Jofré-Saldía E, Torres-Banduc M, Galdames Maliqueo S, Barahona-Fuentes G, Cofré Acevedo C, et al. Effects of a Low Dose of Orally Administered Creatine Monohydrate on Post-Fatigue Muscle Power in Young Soccer Players. *Nutrients* [Internet]. 2024 Apr 28 [cited 2026 May 15];16(9):[11 p.]. Available from: <https://doi.org/10.3390/nu16091324>.
10. Harmancı H, Kalkavan A, Karavelioğlu MB, Şentürk A. Effects of Creatine Supplementation on Motor Performance in Female Futsal Players. *Online J Recreat Sport* [Internet]. 2013 Jan 1 [cited 2026 May 15];2(4):[7 p.]. Available from: <https://tojqi.net/journals/tojras/articles/v02i04/v02i04-04.pdf>.
11. Garcia MP, Longobardi I, Saito T, Miranda MS, Roschel H, Gualano B. Safety of Long-Term Creatine Supplementation in Women’s Football Players: A Real-World In-Season Study. *J Int Soc Sports Nutr* [Internet]. 2025 Sep 30 [cited 2026 May 15];22(Suppl 1):[9 p.]. Available from: <https://doi.org/10.1080/15502783.2025.2591782>.
12. De Zan D, Eletti F, Fiore G, Di Girolamo E, Bozzini GGM, Perico V, et al. Use of Nutritional Strategies, Bioactive Compounds, and Dietary Supplements in Young Athletes:

- From Evidence to Potential Risks-A Narrative Review. *Nutrients* [Internet]. 2025 Jun 30 [cited 2026 May 15];17(13):[23 p.]. Available from: <https://doi.org/10.3390/nu17132194>.
13. North M, Kelly AL, Ranchordas MK, Cole M. Nutritional Considerations in High Performance Youth Soccer: A Systematic Review. *J Sci Sport Exerc*. 2022 Aug;4(3):195–212.
 14. Wang CC, Lin SC, Hsu SC, Yang MT, Chan KH. Effects of Creatine Supplementation on Muscle Strength and Optimal Individual Post-Activation Potentiation Time of the Upper Body in Canoeists. *Nutrients* [Internet]. 2017 Oct 27 [cited 2026 May 15];9(11):[10 p.]. Available from: <https://doi.org/10.3390/nu9111169>.
 15. Wu SH, Chen KL, Hsu C, Chen HC, Chen JY, Yu SY, et al. Creatine Supplementation for Muscle Growth: A Scoping Review of Randomized Clinical Trials from 2012 to 2021. *Nutrients* [Internet]. 2022 Mar 16 [cited 2026 May 15];14(6):[17 p.]. Available from: <https://doi.org/10.3390/nu14061255>.
 16. Du H, Liu S, Li M, Zhao K, Jiang W, You T, et al. Effects of Nutritional Supplements on Explosive Lower Limb Performance in Volleyball Players: A Systematic Review and Network Meta-Analysis. *Nutrients* [Internet]. 2025 Nov 26 [cited 2026 May 15];17(23):[26 p.]. Available from: <https://doi.org/10.3390/nu17233702>.
 17. Atakan MM, Karavelioğlu MB, Harmancı H, Cook M, Bulut S. Short Term Creatine Loading without Weight Gain Improves Sprint, Agility and Leg Strength Performance in Female Futsal Players. *Sci Sports*. 2019 Oct;34(5):321–327.
 18. Valenta S, Psotta R, Hrabal J. The Effects of Creatine Supplementation on Single and Intermittent Anaerobic Exercises and Body Composition During Reduced Training in Soccer Players. *Acta Salus Vitae*. 2022 Jul;10(1):58–71.
 19. Yáñez-Silva A, Buzzachera CF, Piçarro IDC, Januario RSB, Ferreira LHB, McAnulty SR, et al. Effect of Low Dose, Short-Term Creatine Supplementation on Muscle Power Output in Elite Youth Soccer Players. *J Int Soc Sports Nutr* [Internet]. 2017 Feb 7 [cited 2026 May 15];14:[8 p.]. Available from: <https://doi.org/10.1186/s12970-017-0162-2>.
 20. Ashtary-Larky D, Candow DG, Forbes SC, Hajizadeh L, Peacock CA, Kalman D, et al. Creatine Supplementation in Combat Sport Athletes: A Narrative Systematic Review. *J Diet Suppl*. 2025 Aug;22(6):844–869.
 21. Mielgo-Ayuso J, Calleja-Gonzalez J, Marqués-Jiménez D, Caballero-García A, Córdova A, Fernández-Lázaro D. Effects of Creatine Supplementation on Athletic Performance in Soccer Players: A Systematic Review and Meta-Analysis. *Nutrients* [Internet]. 2019 Mar 31 [cited 2026 May 15];11(4):[17 p.]. Available from: <https://doi.org/10.3390/nu11040757>.
 22. Juhász I, Györe I, Csende Z, Rácz L, Tihanyi J. Creatine Supplementation Improves the Anaerobic Performance of Elite Junior Fin Swimmers. *Acta Physiol Hung*. 2009 Sep;96(3):325–336.
 23. Fabiano N, Candow D. Creatine Supplementation: More Is Likely Better for Brain Bioenergetics, Health and Function. *J Psychiatry Brain Sci* [Internet]. 2025 Jul 23 [cited 2026 May 15];10(4):[11 p.]. Available from: <https://doi.org/10.20900/jpbs.20250006>

24. Kious BM, Kondo DG, Renshaw PF. Creatine for the Treatment of Depression. *Biomolecules* [Internet]. 2019 Aug 23 [cited 2026 May 15];9(9):[25 p.]. Available from: <https://doi.org/10.3390/biom9090406>.
25. Prokopidis K, Giannos P, Triantafyllidis KK, Kechagias KS, Forbes SC, Candow DG. Effects of Creatine Supplementation on Memory in Healthy Individuals: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Nutr Rev*. 2023 Mar;81(4):416–427.
26. Moriarty T, Bourbeau K, Dorman K, Runyon L, Glaser N, Brandt J, et al. Dose-Response of Creatine Supplementation on Cognitive Function in Healthy Young Adults. *Brain Sci* [Internet]. 2023 Sep 1 [cited 2026 May 15];13(9):[10 p.]. Available from: <https://doi.org/10.3390/brainsci13091276>.
27. Rawson ES, Lieberman HR, Walsh TM, Zuber SM, Harhart JM, Matthews TC. Creatine Supplementation Does Not Improve Cognitive Function in Young Adults. *Physiol Behav*. 2008 Sep;95(1–2):130–134.
28. Kreider RB, Jäger R, Purpura M. Bioavailability, Efficacy, Safety, and Regulatory Status of Creatine and Related Compounds: A Critical Review. *Nutrients* [Internet]. 2022 Feb 28 [cited 2026 May 15];14(5):[51 p.]. Available from: <https://doi.org/10.3390/nu14051035>.
29. Kreider RB, Jagim AR, Antonio J, Kalman DS, Kerksick CM, Stout JR, et al. Creatine Supplementation Is Safe, Beneficial Throughout the Lifespan, and Should Not Be Restricted. *Front Nutr* [Internet]. 2025 Apr 4;12:[7 p.]. Available from: <https://doi.org/10.3389/fnut.2025.1578564>.
30. de Guingand DL, Palmer KR, Snow RJ, Davies-Tuck ML, Ellery SJ. Risk of Adverse Outcomes in Females Taking Oral Creatine Monohydrate: A Systematic Review and Meta-Analysis. *Nutrients* [Internet]. 2020 Jun 15 [cited 2026 May 15];12(6):[26 p.]. Available from: <https://doi.org/10.3390/nu12061780>.
31. Gualano B, Ugrinowitsch C, Novaes RB, Artioli GG, Shimizu MH, Seguro AC, et al. Effects of Creatine Supplementation on Renal Function: A Randomized, Double-Blind, Placebo-Controlled Clinical Trial. *Eur J Appl Physiol*. 2008 May;103(1):33–40.
32. Everett S. Optimizing Performance Nutrition for Adolescent Athletes: A Review of Dietary Needs, Risks, and Practical Strategies. *Nutrients* [Internet]. 2025 Aug 28 [cited 2026 May 15];17(17):[20 p.]. Available from: <https://doi.org/10.3390/nu17172792>
33. American Dietetic Association, Dietitians of Canada, American College of Sports Medicine, Rodriguez NR, Di Marco NM, Langley S. American College of Sports Medicine Position Stand. Nutrition and Athletic Performance. *Med Sci Sports Exerc*. 2009 Mar;41(3):709–731.
34. Pakulak A, Candow DG, Totony de Zepetnek J, Forbes SC, Basta D. Effects of Creatine and Caffeine Supplementation During Resistance Training on Body Composition, Strength, Endurance, Rating of Perceived Exertion and Fatigue in Trained Young Adults. *J Diet Suppl*. 2022 Mar;19(5):587–602.

35. Jagim AR, Stecker RA, Harty PS, Erickson JL, Kerksick CM. Safety of Creatine Supplementation in Active Adolescents and Youth: A Brief Review. *Front Nutr* [Internet]. 2018 Nov 28 [cited 2026 May 15];5:[13 p.]. Available from: <https://doi.org/10.3389/fnut.2018.00115>.
36. Filler G, Maung E, Díaz González de Ferris ME, Chan NG, Sharma AP. Acute Kidney Injury with Cast Nephropathy Following Creatine Loading in a 17-Year-Old. *Pediatr Nephrol*. 2025 Oct;40(10):3089–3092.
37. Davani-Davari D, Karimzadeh I, Ezzatzadegan-Jahromi S, Sagheb MM. Potential Adverse Effects of Creatine Supplement on the Kidney in Athletes and Bodybuilders. *Iran J Kidney Dis*. 2018 Oct;12(5):253–260.
38. Simpson AJ, Horne S, Sharp P, Sharps R, Kippelen P. Effect of Creatine Supplementation on the Airways of Youth Elite Soccer Players. *Med Sci Sports Exerc*. 2019 Aug;51(8):1582–1590.
39. van der Merwe J, Brooks NE, Myburgh KH. Three Weeks of Creatine Monohydrate Supplementation Affects Dihydrotestosterone to Testosterone Ratio in College-Aged Rugby Players. *Clin J Sport Med*. 2009 Sep;19(5):399–404.
40. Silva C, Fung AWS, Masson V, Assen K, Ward V, McKenzie J, et al. Vitamin D Toxicity from an Unusual and Unexpected Source: A Report of 2 Cases. *Horm Res Paediatr*. 2023 Jun;96(3):332–340.
41. LaBotz M, Griesemer BA. Use of Performance-Enhancing Substances. *Pediatrics* [Internet]. 2016 Jul 1 [cited 2026 May 15];138(1):[15 p.]. Available from: <https://doi.org/10.1542/peds.2016-1300>.
42. Ganson KT, Mitchison D, Murray SB, Nagata JM. Legal Performance-Enhancing Substances and Substance Use Problems Among Young Adults. *Pediatrics* [Internet]. 2020 Sep 1 [cited 2026 May 15];146(3):[9 p.]. Available from: <https://doi.org/10.1542/peds.2020-0409>.
43. Jagim AR, Kerksick CM. Creatine Supplementation in Children and Adolescents. *Nutrients* [Internet]. 2021 Feb 18 [cited 2026 May 15];13(2):[17 p.]. Available from: <https://doi.org/10.3390/nu13020664>.
44. Metzger GA, Minneci PM, Gehred A, Day A, Klingele KE. Creatine Supplementation in the Pediatric and Adolescent Athlete--A Literature Review. *J Orthop*. 2023 Apr;38(3):73–78.
45. Hall M, Trojjan TH. Creatine Supplementation. *Curr Sports Med Rep*. 2013 Jul-Aug;12(4):240–244.

Assessing Adherence for Counselling Non–Insulin-Dependent Type 2 Diabetics Regarding “Sick-Day Rules”: A Chart Review of 300 Chronic Disease Management (CDM) Patients at the Family Medicine Unit in Regina, Saskatchewan

Manraj Aujla, FMR II; Manav Bhalla, FMR II; Sanchsith Rajalingam, FMR II
Rejina Kamrul, MD, CCFP; Adam Clay, MSc

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Older adults with type 2 diabetes mellitus (T2DM) often face increased risks of dehydration, hypoglycemia, acute kidney injury, and hospitalization during illness due to multiple comorbidities and medications. “Sick-day rules” advise temporary medication adjustments to prevent such complications but are inconsistently applied in primary care.

Question(s): What is the proportion of patients with non–insulin-dependent T2DM at the Family Medicine Unit in Regina, Saskatchewan, who received counselling and written reference materials regarding sick-day rules?

Methods/Methodology: Patients aged 65 years or older with non–insulin-dependent T2DM who were prescribed at least one glucose-lowering medication were identified using the reporting function in the Electronic Medical Record (EMR). Patients were enrolled in the CDM roster at the Family Medicine Unit for at least 12 months. Records were then manually reviewed to collect demographics, comorbidities (congestive heart failure, chronic kidney disease, and coronary artery disease), and documentation of counselling or written resources provided during clinic visits. Descriptive statistics and Fisher’s exact tests were conducted. Ethics Approval was received from the University of Saskatchewan’s Biomedical REB (Bio ID 5831) and Operational approval (OA-Bio-5831) was granted by the Saskatchewan Health Authority.

Results/Findings: The cohort captured in our study (n=299) had a median age of 73 years (IQR, 69–79), and 48.7% (n=146) were male. Overall, 31.3% (n=94) had documented sick-day counselling by a physician at the Family Medicine Unit, and 19.7% (n=59) had documentation of written materials provided during the encounter. Counselling rates did not differ significantly by age, sex, or comorbidity status (p>0.05). Of the 206 patients without documented counselling, 63 had major comorbidities.

Discussion: Documentation of sick-day education was suboptimal, particularly for patients with significant comorbidities. The findings are consistent with current literature, which demonstrates inconsistent clinician familiarity, a lack of standardized workflows, and limited EMR prompts to support documentation.

Conclusions: Sick-day counselling and written resource provision were infrequent and inconsistent, highlighting a significant learning opportunity for physicians managing patients with T2DM.

Recommendations: Integrating EMR prompts, easy-access patient handouts, and targeted clinician education may improve consistency of counselling efforts. Implementing these supports, along with a follow-up audit, is recommended to assess their impact.

References:

1. Wu Y, Ding Y, Tanaka Y, Zhang W. Risk Factors Contributing to Type 2 Diabetes and Recent Advances in Treatment and Prevention. *Int J Med Sci*. 2014 Sep;11(11):1185–1200.
2. Dhaliwal KK, Watson KE, Lamont NC, Drall KM, Donald M, James MT, et al. Managing “Sick Days” in Patients with Chronic Conditions: An Exploration of Patient and Healthcare Provider Experiences. *Health Expect*. 2023 Aug;26(4):1746-1756.
3. Kado K, Okada H, Suzuki S, Satake M, Yamazaki T, Kurosawa M, et al. Study of Assessment of Knowledge and Understanding for Coping with Sick Days Among Patients with Diabetes in Community Pharmacy: A Cluster Randomized Controlled Trial (SAKURA Trial). *J Pharm Policy Pract* [Internet]. 2023 Sep 25 [cited 2026 May 13];16(1):[10 p.]. Available from: <https://doi.org/10.1186/s40545-023-00614-4>.
4. Watson KE, Dhaliwal K, McMurtry E, Donald T, Lamont N, Benterud E, et al. Sick Day Medication Guidance for People with Diabetes, Kidney Disease, or Cardiovascular Disease: A Systematic Scoping Review. *Kidney Med* [Internet]. 2022 May 28 [cited 2026 May 13];4(9):[12 p.]. Available from: <https://doi.org/10.1016/j.xkme.2022.100491>.
5. Berg EG. Sick-Day Rules: What to Do When You Are Feeling Under the Weather. *Diabetes Forecast*. 2013 Oct;66(10):32–33.
6. Whiting P, Morden A, Tomlinson LA, Caskey F, Blakeman T, Tomson C, et al. What Are the Risks and Benefits of Temporarily Discontinuing Medications to Prevent Acute Kidney Injury? A Systematic Review and Meta-Analysis. *BMJ Open* [Internet]. 2017 Apr 7 [cited 2026 May 13];7(4):[9 p.]. Available from: <https://doi.org/10.1136/bmjopen-2016-012674>.
7. Watson KE, Dhaliwal K, Robertshaw S, Verdin N, Benterud E, Lamont N, et al. Consensus Recommendations for Sick Day Medication Guidance for People with Diabetes, Kidney, or Cardiovascular Disease. *Am J Kidney Dis*. 2023 May;81(5):564-574.
8. Yuang K, Espinoza AL, McCoy RG. Glycemic Management in Adults with Diabetes During Illness. *Clin Diabetes*. 2025 Sep;43(5):735-745.
9. Endotext [Internet]. Bethesda (MD): National Library of Medicine; c2026. Dietary advice for individuals with diabetes; 2024 Apr 28 [cited 2026 May 13];[about 28 screens]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK279012/>.

Appropriate primary care prescribing and patient compliance with compression therapy for venous insufficiency

Maham Tasneem, FMR II; Warda Zanib, FMR II; Adam Clay, MSc
Rejina Kamrul, MBBS, CCFP; Abdalla Butt, MD, FRCPC; David Kopriva, MD, CM, FRCPC.

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Varicose veins and chronic venous insufficiency are common conditions that may significantly impact patient quality of life. Most cases can be effectively managed with compression therapy administered in primary care settings.

Question(s): In patients referred to vascular surgery for varicose veins or venous insufficiency, how often do family physicians prescribe compression therapy prior to referral, and what is the rate of patient adherence to prescribed compression therapy?

Methods/Methodology: A retrospective chart review was completed for patients of family physicians working at the Family Medicine Unit, The Nest, Greens Medical Clinic, Stapleford Medical Clinic or Victoria East Medical Clinic. Patients referred to CVT Associates were identified using the CVT Associates database. Vascular surgery consult notes and family physician referral letters were reviewed to determine whether compression therapy was prescribed before referral and whether compliance was documented. This project was approved by the University of Saskatchewan's Biomedical Research Ethics Board (Bio 6047).

Results/Findings: A total of 287 patients were identified based on the CVT Associates database. Manual chart review identified 113 patients referred for varicose veins/venous insufficiency. Referrals per family physician ranged from 1 to 10. Before January 31, 2021, 19 of 76 referrals (25.0%) documented compression therapy prescription. After January 31, 2021, 12 of 37 referrals (32.4%) documented compression therapy prescription. Overall, 31 of 113 referrals (27.4%) documented compression therapy before referral. Patient compliance was noted at CVT in 10 of 76 patients (13.2%) before January 31, 2021, and 10 of 35 patients seen after January 31, 2021 (28.6%). Most patients were not offered intervention due to a lack of indications for invasive therapy.

Discussion: These findings suggest documentation of compression therapy prior to referral is often inconsistent, highlighting an opportunity for education within the primary care setting.

Conclusions: Compression therapy was documented in a minority of referrals, despite being first line conservative management.

Recommendations: Future improvement could include teaching, clearer documentation of compression therapy education, and referral prompts encouraging a trial of conservative management before vascular surgery referral.

References:

1. Lim CS, Davies AH. Graduated Compression Stockings. *CMAJ*. 2014 Jul;186(10):E391-398.
2. Knight SL, Robertson L, Stewart M. Graduated Compression Stockings for the Initial Treatment of Varicose Veins in People without Venous Ulceration. *Cochrane Database Syst Rev* [Internet]. 2021 Jul 16 [cited 2026 May 15];7(7):[71 p.]. Available from: <https://doi.org/10.1002/14651858.cd008819.pub4>.
3. Sell H, Vikatmaa P, Albäck A, Lepäntalo M, Malmivaara A, Mahmoud O, et al. Compression Therapy Versus Surgery in the Treatment of Patients with Varicose Veins: A RCT. *Eur J Vasc Endovasc Surg*. 2014 Jun;47(6):670-677.
4. Mestre S, Triboulet J, Demattei C, Veye F, Nou M, Pérez-Martin A, et al. Acute Effects of Graduated and Progressive Compression Stockings on Leg Vein Cross-Sectional Area and Viscoelasticity in Patients with Chronic Venous Disease. *J Vasc Surg Venous Lymphat Disord*. 2022 Jan;10(1):186-195.
5. Tan M, Urbanek T, Rabe E, Giancesini S, Parsi K, Davies AH. Compression Therapy in the Management of Varicose Veins. *Phlebology*. 2024 May;39(4):276-279.
6. Attaran RR, Edwards ML, Arena FJ, Bunte MC, Carr JG, Castro-Dominguez Y, et al. 2025 SCAI Clinical Practice Guidelines for the Management of Chronic Venous Disease: This Statement Was Endorsed by the Society for Vascular Medicine (SVM). *J Soc Cardiovasc Angiogr Interv* [Internet]. 2025 Aug 30 [cited 2026 May 15];4(8):[15 p.]. Available from: <https://doi.org/10.1016/j.jscai.2025.103729>

Evaluating Physicians' Perceptions of Procedural Skills Taught in Saskatchewan's Family Medicine Surgical Skills Curriculum

Chaudhry Masud, FMR II; Husam Muftah, FMR II; Jack Yassa, FMR II;
Clara Rocha Michaels, MD, CCFP; Adam Clay, MSc;
Andrea Vasquez Camargo, MD, MSc, CCFP.

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Procedural competence is an essential component of comprehensive family medicine training. However, family physicians may perform fewer procedures after residency due to limited confidence, low clinical volume, or practice environment constraints. Understanding graduates' perceptions of the Family Medicine Surgical Skills Curriculums in Saskatchewan can identify whether training is relevant to independent practice and where curriculum improvements are needed.

Question(s): How do University of Saskatchewan Family Medicine graduates perceive the relevance, confidence, and applicability of procedural skills taught during residency? What barriers to procedural practice and curriculum improvements do they identify?

Methods/Methodology: A cross-sectional online survey with closed- and open-ended items was distributed to family physicians who graduated from Saskatchewan's Family Medicine Program between July 1, 2017 and June 30, 2024. Quantitative data were summarized using descriptive statistics, and data from open-ended responses were reviewed to identify recurring themes. Ethics approval was obtained through the University of Saskatchewan Behavioural Research Ethics Board (Beh 6177).

Results/Findings: 34 respondents were analyzed (34/312 = 10.9%), including urban 19 (55.9%), regional 13 (38.2%), and rural 2 (5.9%) physicians. Twenty-one (61.7%) graduated within 5 years. Most graduates (33–34; 97–100%) intended to perform local anesthetic and skin procedures; only 4–12 respondents (11.7–35.2%) were not currently performing them. Fewer than half performed most gastrointestinal, genitourinary, and MSK procedures, except shoulder 25 (78.1%) and knee injections 28 (84.8%). Practice environment influenced expected procedural use for 29/31 respondents (93.5%). Barriers included limited time, equipment availability, lack of confidence or experience, and limited opportunities. Suggested improvements included more hands-on workshops, simulation-based training, mentorship, greater procedural volume, and repeated exposure.

Discussion: Findings align with literature showing that structured curricula, simulation, mentorship, and repeated exposure support procedural confidence, while limited clinical volume remains a barrier. Although respondents perceived the curriculums as relevant, variable confidence and use suggest that ongoing procedural opportunities are needed.

Conclusions: The Surgical Skills Curriculums in Saskatchewan appears relevant to family medicine practice, but opportunities exist to strengthen confidence and sustain procedural use after graduation.

Recommendations: Future curriculum development should prioritize repeated hands-on exposure, simulation, mentorship, and structured opportunities across both urban and rural sites.

References:

1. Sy J, Bloom J, Mizdrak N, Moaveni A, Toubassi D. Procedural Skills Training During Residency: Introduction of a Procedure Clinic to Improve Technical Competencies. *Can Fam Physician*. 2023 Apr;69(4):292-294.
2. Bansal S, Bhambhwani V. Evaluation of a Simulation-Based Ophthalmology Education Workshop for Medical Students: A Pilot Project. *BMC Med Educ* [Internet]. 2025 Jan 30 [cited 2026 May 14];25(1):[8 p.]. Available from: <https://doi.org/10.1186/s12909-025-06712-y>.
3. Branditz LD, Kendle AP, Leung CG, San Miguel CE, Way DP, Panchal AR, Yee J. Bridging the Procedures Skill Gap from Medical School to Residency: A Simulation-Based Mastery Learning Curriculum. *Med Educ Online* [Internet]. 2024 Dec 31 [cited 2026 May 14];29(1):[7 p.]. Available from: <https://doi.org/10.1080/10872981.2024.2412399>.
4. Chatterji R, Foote J, Fry M, Erwin A, Crutcher J, Kesto W. Simulated Subacromial Injection Instruction Improves Accuracy and Skill Level: A Model for Musculoskeletal Procedural Training. *BMC Med Educ* [Internet]. 2024 May 14 [cited 2026 May 14];24(1):[9 p.]. Available from: <https://doi.org/10.1186/s12909-024-05456-5>.
5. Clemens L. The Efficacy and Cost-Effectiveness of a Simulation-Based Primary Care Procedural Skills Training Program for Advanced Practice Providers. *J Contin Educ Health Prof*. 2024 Jul;44(3):222-228.
6. Rezaiefar P, Archibald D, Kabir M, Humphrey-Murto S. Challenges in Providing Gynecological Procedures in Primary Care: A Survey of Canadian Academic Family Physicians. *Womens Health Rep (New Rochelle)*. 2025 Jan;6(1):102-112.
7. Khan WU, Twomey J, Ryan E, Martin T, Kamal M, Lok Boris Cheng P, et al. Barriers and Enablers to Achieving Clinical Procedure Competency-Based Outcomes in a National Paediatric Training/Residency Program-A Multi-Centered Qualitative Study. *BMC Med Educ* [Internet]. 2023 Dec 13 [2026 May 14];23(1):[10 p.]. Available from: <https://doi.org/10.1186/s12909-023-04928-4>.

A Rapid Review of Principles, Practices, and Challenges in Canada, Australia, and New Zealand Regarding the Teaching of Indigenous Health in Medical Education

Hamza Dani, FMR II; Aaron Prystupa, MD, CCFP
Vivian R Ramsden, RN, PhD, MCFP (Hon.), FCAHS; R Bryce, MD, MSc

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Indigenous populations in Canada continue to experience significant health inequities rooted in colonization and systemic racism. Medical education institutions are at the forefront of reconciliation and are key organizations in addressing the Truth and Reconciliation Commission of Canada's Call to Action #24 regarding Indigenous health education for health professionals.

Question(s):

1. What principles, strategies, and exemplary practices have other institutions both in Canada and Australia/New Zealand utilized to promote successful integration of Indigenous health curriculum into their medical training programs?
2. What challenges to the integration of Indigenous health curriculum into medical training programs have been identified in Canadian and Australia/New Zealand contexts?

Methods/Methodology: A rapid review of the PubMed database was commenced on September 15, 2025. Following structured screening of 238 articles by two reviewers using predefined inclusion and exclusion criteria, 32 studies were included and analyzed using thematic analysis. Due to the presence of all study materials in the public domain, formal ethics approval was not required.

Results/Findings: Successful Indigenous health harmonization is characterized by community-led partnerships, longitudinal immersion, and a collaborative approach. However, significant barriers remain, including the persistence of institutional racism, resource scarcity, and already overloaded curriculum.

Discussion: This review highlights that effective Indigenous health education extends beyond content delivery to relational, structural, and experiential learning. Community-led partnerships and immersive experiences are central to fostering cultural humility and advocacy. However, their implementation is constrained by limited resources, competing curricular demands, and insufficient institutional commitment. The findings show a gap between policy expectations and practical execution, suggesting that without sustained structural support and accountability, efforts risk remaining superficial rather than transformative.

Conclusions: Effective harmonization requires moving toward a model that emphasizes lifelong self-critique and institutional support.

Recommendations: Future reforms must focus on building partnerships to ensure Indigenous health is a core, rather than marginal, component of medical professionalism, with an emphasis on community-led, immersive, collaborative models.

References:

1. Bethune R, Absher N, Obiagwu M, Qarmout T, Steeves M, Yaghoubi M, et al. Social Determinants of Self-Reported Health for Canada’s Indigenous Peoples: A Public Health Approach. *Public Health*. 2019 Nov;176(11):172–180.
2. Joseph G. The Role of Sovereignty in Indigenous Community-Based Health Interventions: A Qualitative Metasynthesis. *Am J Community Psychol*. 2024 Mar;73(1–2):216–233.
3. Reading C, Wein F. Health Inequalities and Social Determinants of Aboriginal Peoples’ Health [Internet]. Prince George (BC): National Collaborating Centre for Aboriginal Health; 2009 [cited 2026 May 15]. 36 p. Available from: <https://www.ccnsa-nccah.ca/docs/determinants/RPT-HealthInequalities-Reading-Wien-EN.pdf>.
4. Allan B, Smylie J. First Peoples, Second Class Treatment: The Role of Racism in the Health and Well-Being of Indigenous Peoples in Canada [Internet]. Toronto (ON): The Wellesley Institute; 2015 [cited 2026 May 10]. 71 p. Available from: <https://www.wellesleyinstitute.com/wp-content/uploads/2015/02/Report-First-Peoples-Second-Class-Treatment-Final.pdf>.
5. National Collaborating Centre for Indigenous Health. Access to Health Services as a Social Determinant of First Nations, Inuit and Métis Health [Internet]. Prince George (BC): National Collaborating Centre for Indigenous Health; 2019 [cited 2026 May 10]. 16 p. Available from: <https://www.nccih.ca/docs/determinants/FS-AccessHealthServicesSDOH-2019-EN.pdf>.
6. Sauvé A, Cappelletti A, Murji L. Stand Up for Indigenous Health: A Simulation to Educate Residents About the Social Determinants of Health Faced by Indigenous Peoples in Canada. *Acad Med*. 2022 Apr;97(4):518–523.
7. National Centre for Truth and Reconciliation [Internet]. Winnipeg (MB): University of Manitoba; c2026. Truth and Reconciliation Commission of Canada; [cited 2026 May 15];[about 5 screens]. Available from: <https://nctr.ca/about/truth-and-reconciliation-commission-of-canada/>.
8. Truth and Reconciliation Commission of Canada. Truth and Reconciliation Commission of Canada: Calls to Action [Internet]. Winnipeg (MB): Truth and Reconciliation Commission of Canada; 2015 [cited 2026 May 15]. 20 p. Available from: https://ehprnh2mwo3.exactdn.com/wp-content/uploads/2021/01/Calls_to_Action_English2.pdf.
9. Curtis E, Loring B, Jones R, Tipene-Leach D, Walker C, Paine SJ, et al. Refining the Definitions of Cultural Safety, Cultural Competency and Indigenous Health: Lessons from Aotearoa New Zealand. *Int J Equity Health* [Internet]. 2025 May 9 [cited 2026 May 15];24(1):[10 p.]. Available from: <https://doi.org/10.1186/s12939-025-02478-3>

10. So N, Price K, O'Mara P, Rodrigues MA. The Importance of Cultural Humility and Cultural Safety in Health Care. *Med J Aust*. 2024 Jan;220(1):12–13.
11. Kurtz DLM, Janke R, Vinek J, Wells T, Hutchinson P, Froste A. Health Sciences Cultural Safety Education in Australia, Canada, New Zealand, and the United States: A Literature Review. *Int J Med Educ*. 2018 Oct;9(1):271–285.
12. Rashid M, Clark W, Foulds JL, Nguyen J, John I, Forgie SE. Exploring Approaches to Teaching Indigenous Health Curricula from the Perspectives of Faculty and Residents. *Med Educ Online* [Internet]. 2025 Dec 31 [cited 2026 May 15];30(1):[9 p.]. Available from: <https://doi.org/10.1080/10872981.2025.2516673>.
13. Jacklin K, Strasser R, Peltier I. From the Community to the Classroom: The Aboriginal Health Curriculum at the Northern Ontario School of Medicine. *Can J Rural Med*. 2014 Fall;19(4):143–150.
14. Zaretsky L, Crooks R, Whalen-Browne M, Gausvik AL, Roach P. Evaluation of a Longitudinal Indigenous Health Elective in Family Medicine. *Can Med Educ J*. 2025 Feb;16(1):65-70.
15. Huria T, Palmer S, Beckert L, Lacey C, Pitama S. Indigenous Health: Designing a Clinical Orientation Program Valued by Learners. *BMC Med Educ* [Internet]. 2017 Oct 5 [cited 2026 May 15];17(1):[8 p.]. Available from: <https://doi.org/10.1186/s12909-017-1019-8>
16. Jarvis-Selinger S, Ho K, Novak Lauscher H, Liman Y, Stacy E, Woollard R, et al. Social Accountability in Action: University-Community Collaboration in the Development of an Interprofessional Aboriginal Health Elective. *J Interprof Care*. 2008 Jan;22(Suppl 1):61–72.
17. Gallagher P, Mckinlay E, Pullon S, McHugh P. Student Perceptions of Cultural Immersion During an Interprofessional Programme. *J Interprof Care*. 2019 Mar-Apr;33(2):264–266.
18. Lacey C, Huria T, Beckert L, Gilles M, Pitama S. The Hui Process: A Framework to Enhance the Doctor–Patient Relationship with Maori. *N Z Med J*. 2011 Dec;124(1347):72-78.
19. Harrison RL, MacNab AJ, Duffy DJ, Benton DHJ. Brighter Smiles: Service Learning, Inter-Professional Collaboration and Health Promotion in a First Nations Community. *Can J Public Health*. 2006 May-Jun;97(3):237–240.
20. Askew DA, Lyall VJ, Ewen SC, Paul D, Wheeler M. Understanding Practitioner Professionalism in Aboriginal and Torres Strait Islander Health: Lessons from Student and Registrar Placements at an Urban Aboriginal and Torres Strait Islander Primary Healthcare Service. *Aust J Prim Health*. 2017 Oct;23(5):446–450.
21. Hall K, Vervoort S, Del Fabbro L, Rowe Minniss F, Saunders V, Martin K, et al. Evolving Beyond Antiracism: Reflections on the Experience of Developing a Cultural Safety Curriculum in a Tertiary Education Setting. *Nurs Inq* [Internet]. 2022 Sep 9 [cited 2026 May 15];30(1):[11 p.]. Available from: <https://doi.org/10.1111/nin.12524>.
22. Paul D. Creating Change: Building the Capacity of the Medical Workforce in Aboriginal Health. *ANZ J Surg*. 2013 Jan;83(1–2):55–59.

23. Sen Gupta TK, Muray RB, McDonell A, Murphy B, Underhill AD. Rural Internships for Final Year Students: Clinical Experience, Education and Workforce. *Rural Remote Health* [Internet]. 2008 Feb 13 [cited 2026 May 15];8(1):[10 p.]. Available from: <https://www.rrh.org.au/journal/article/827>.
24. Jumah NA, Wilson D, Shah R. A Canadian Survey of Postgraduate Education in Aboriginal Women's Health in Obstetrics and Gynaecology. *J Obstet Gynaecol Can*. 2013 Jul;35(7):647–653.
25. Vass A, Adams K. Educator Perceptions on Teaching Indigenous Health: Racism, Privilege and Self-Reflexivity. *Med Educ*. 2021 Feb;55(2):213–221.
26. Rashid M, Clark W, Foulds JL, John I, Nguyen J, Chan M, et al. Indigenous Elders and Academic Leaders' Perspectives on Training Residents to Work with Indigenous Populations: A Constructivist Grounded Theory Study. *Clin Teach* [Internet]. 2025 Sep 29 [cited 2026 May 15];22(6):[13 p.]. Available from: <https://doi.org/10.1111/tct.70219>
27. Manton D, Williams M, Hayen A. "It's About Rights": The Bunya Project's Indigenous Australian Voices on Health Care Curricula and Practice. *Health Hum Rights*. 2024 Jun;26(1):87–100.
28. Rashid M, Nguyen J, Clark W, Foulds JL, John I, Chan MK, et al. Residency Training Programs to Support Residents Working in First Nations, Inuit, and Métis Communities. *BMC Med Educ* [Internet]. 2025 Jan 28 [cited 2026 May 15];25(1):[14 p.]. Available from: <https://doi.org/10.1186/s12909-025-06722-w>.
29. Riediger ND, Cyr M, Mignone J. An Evaluation of an Experiential Learning Program in Global and Indigenous Health: The University of Manitoba's Queen Elizabeth II Diamond Jubilee Scholarship Program. *Inquiry* [Internet]. 2020 Aug 26 [cited 2026 May 15];57:[10 p.]. Available from: <https://doi.org/10.1177/0046958020951002>.
30. Durie M. Indigenizing Mental Health Services: New Zealand Experience. *Transcult Psychiatry*. 2011 Apr;48(1–2):24–36.
31. Pinnock R, Jones R, Wearn A. Learning and Assessing Cultural Competence in Paediatrics. *Med Educ*. 2008 Nov;42(11):1124–1125.
32. Jones RG, Henning MA, Pinnock R, Shulruf B, Hawken J. Medical Students' and Clinical Teachers' Perceptions of Maori Health Teaching. *N Z Med J*. 2013 Jun;126(1377):40–49.
33. Watt K, Abbott P, Reath J. Cultural Competency Training of GP Registrars-Exploring the Views of GP Supervisors. *Int J Equity Health* [Internet]. 2015 Oct 6 [cited 2026 May 15];14(1):[10 p.]. Available from: <https://doi.org/10.1186/s12939-015-0226-3>.
34. Yeung S, Bombay A, Walker C, Denis J, Martin D, Sylvestre P, et al. Predictors of Medical Student Interest in Indigenous Health Learning and Clinical Practice: A Canadian Case Study. *BMC Med Educ* [Internet]. 2018 Dec 14 [cited 2026 May 15];18(1):[11 p.]. Available from: <https://doi.org/10.1186/s12909-018-1401-1>.
35. Micheal S, Ogbeide AE, Arora A, Alford S, Firdaus R, Lim D, et al. Exploring Tertiary Health Science Student Willingness or Resistance to Cultural Competency and Safety

- Pedagogy. *Int J Environ Res Public Health* [Internet]. 2021 Aug 31 [cited 2026 May 15];18(17):[14 p.]. Available from: <https://doi.org/10.3390/ijerph18179184>.
36. Larson B, Herx L, Williamson T, Crowshoe L. Beyond the Barriers: Family Medicine Residents' Attitudes Towards Providing Aboriginal Health Care. *Med Educ*. 2011 Apr;45(4):400–406.
 37. Duke M, Ewen S. Implementation of Indigenous Mental Health Training in Victoria. *Australas Psychiatry*. 2009 Jun;17(3):228–232.
 38. Martin ME, Reath JS. General Practice Training in Aboriginal and Torres Strait Islander Health. *Med J Aust*. 2011 Jun;194(11):S67-70.

A Review of Preventative Care in HIV Care

Morgan Schatz, FMR II; Jelyssa Luc, FMR II;
Cole Picot, MD, CCFP; Katrina Sawchuk, MEd, PhD

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Screening for common non-infectious comorbidities is critical to providing comprehensive care for people living with HIV (PLWHIV). West Winds Primary Health Centre (WWPHC) offers outpatient HIV care as part of the Positive Living Program (PLP) services offered by the SHA. This study references the Primary Care Guidelines for the Management of Adults Living With HIV/AIDS in British Columbia (BC-CfE HIV/AIDS Guidelines) to examine the current rate of non-infectious comorbidity screening for PLWHIV at WWPHC.

Question(s): Are PLWHIV who receive care at WWPHC screened for common non-infectious comorbidities in accordance with recommendations set out by the BC-CfE HIV/AIDS Guidelines?

Methods/Methodology: This was a retrospective chart review of 49 patients who are a part of the PLP at WWPHC from January of 2024 to December of 2025. Charts containing a diagnostic code for HIV were de-identified and screening was coded as not up to date, up to date, or missing. Descriptive analyses were performed with the support of CRSU. This project was Exempt from Ethics (E-Bio-090) and received SHA Operational Approval.

Results/Findings: Liver function, renal function, lipid levels, and diabetes were recorded as the screening most often up to date (76.6%-83.7%). Screening investigations such as blood pressure reading that required an in-person appointment was lower at 71.4%. Cervical cancer was the least likely completed screen at 39%.

Discussion: In-office testing such as blood pressure had the highest screening rates followed by serum laboratory testing. Pap testing and FIT had the lowest up to date rates, likely because they are more invasive and difficult to arrange. A lower mean of attended visits compared to scheduled visits, which may indicate larger potential systemic barriers to obtaining healthcare.

Conclusions: This study has identified that there is a gap in screening for non-infectious comorbidities in PLWHIV who receive care at WWPHC.

Recommendations: Consider qualitative studies to identify potential barriers to providing the recommended screening investigations. Multidisciplinary teams and integrated care networks could help identify shared goals and help manage concurrent needs to help optimize and streamline care.

References:

1. CATIE – Canada’s Source for HIV and Hepatitis C Information [Internet]. Toronto (ON): CATIE; c2026. The epidemiology of HIV in Canada; 2026 [cited 2026 May 2];[about 14 screens]. Available from: <https://www.catie.ca/the-epidemiology-of-hiv-in-canada>.
2. Government of Canada [Internet]. Ottawa (ON): Public Health Agency of Canada; c2026. HIV in Canada: 2024 surveillance highlights; [modified 2026 Apr 16; cited 2026 May 5];[about 6 screens]. Available from: <https://www.canada.ca/en/public-health/services/diseases/hiv-aids/health-professionals/surveillance-highlights.html>.
3. Government of Canada [Internet]. Ottawa (ON): Public Health Agency of Canada; c2026. HIV in Canada, Surveillance Report to December 31, 2023: Executive summary; [modified 2025 Sep 18; cited 2026 May 5];[about 8 screens]. Available from: <https://www.canada.ca/en/public-health/services/publications/diseases-conditions/hiv-canada-surveillance-report-december-31-2023.html>.
4. Heron JE, Norman SM, Yoo J, Lembke K, O’Connor CC, Weston CE, et al. The Prevalence and Risk of Non-Infectious Comorbidities in HIV-Infected and Non-HIV Infected Men Attending General Practice in Australia. PLoS One [Internet]. 2019 Oct 9 [cited 2026 May 12];14(10):[13 p.]. Available from: <https://doi.org/10.1371/journal.pone.0223224>.
5. Horberg M, Thompson M, Agwu A, Colasanti J, Haddad M, Jain M, et al. Primary Care Guidance for Providers Who Care for Persons with Human Immunodeficiency Virus: 2024 Update by the HIV Medicine Association of the Infectious Diseases Society of America. Primary Care Guidance for Providers Who Care for Persons with Human Immunodeficiency Virus: 2024 Update by the HIV Medicine Association of the Infectious Diseases Society of America. Clin Infect Dis [Internet]. 2024 Oct 12 [cited 2026 May 12];[about 57 p.]. Available from: <https://doi.org/10.1093/cid/ciae479>. Epub ahead of print. PubMed PMID: 39393187.
6. Shah ASV, Stelzle D, Lee KK, Beck EJ, Alam S, Clifford S, et al. Global Burden of Atherosclerotic Cardiovascular Disease in People Living with HIV: Systematic Review and Meta-Analysis. Circulation. 2018 Sept;138(11):1100–12.
7. British Columbia Centre of Excellence in HIV/AIDS. Primary care guidelines for the management of adults living with HIV/AIDS in British Columbia [Internet]. Vancouver (BC): British Columbia Centre of Excellence; 2011 Mar [updated 2021 Aug; cited 2026 May 12]. 127 p. Available from: <https://bccfe.ca/therapeutic-guidelines/primary-care-guidelines/>.
8. Avari P, Devendra S. Human Immunodeficiency Virus and Type 2 Diabetes. London J Prim Care. 2017 Mar;9(3):38–42.
9. Goh SSL, Lai PS, Tan AT, Ponnampalavanar S. Reduced Bone Mineral Density in Human Immunodeficiency Virus-Infected Individuals: A Meta-Analysis of Its Prevalence and Risk Factors. Osteoporosis Int. 2017 Nov;29(3):595–613.
10. Hernández-Ramírez RU, Shiels MS, Dubrow R, Engels EA. Cancer Risk in HIV-Infected People in the USA from 1996 to 2012: A Population-Based, Registry-Linkage Study. Lancet HIV. 2017 Nov;4(11):e495-504.
11. StatPearls [Internet]. Bethesda (MA): National Library of Medicine; c2026. Evidence-based medicine; 2024 Sep 10 [cited 2026 May 3];[about 5 screens]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK470182/>.
12. Microsoft 365 Copilot. Version bizchat.20260430.50.2 [software]. 2026 Apr 30 [cited 2026 May 12].

Accessing Healthcare and Affirming Care: Perspectives of 2SLGBTQIA+ Individuals in North Battleford, Saskatchewan

Jessica Kainth, FMR II; Sarah Nazim, FMR I; Michael Barnett MD, CCFP
Ginger Ruddy, MD, CCFP; Rhonda Bryce, MD, MSc

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Family medicine aims to provide inclusive and accessible care to everyone, including members of the 2SLGBTQIA+ community who may not feel comfortable presenting for care. With support from local organizations, we hoped to better understand the experiences and perspectives of community members to suggest improvements to North Battleford's primary care, public health, and urgent/emergent care services.

Question(s): This project aims to examine the quality, accessibility, and potential improvement of front-line services for 2SLGBTQIA+-identifying patients in North Battleford.

Methods/Methodology: Distributed through Battlefords Pride and Battle River Treaty 6 Health Centre, this cross-sectional, mixed qualitative/quantitative online survey for 2SLGBTQIA+-identifying individuals inquired about experiences related to primary care, public health clinics, walk-in clinics, or emergency department services in North Battleford, Saskatchewan during the last five years. This study was exempted from Research Ethics Board review (E-Bio-129) as quality improvement work.

Results/Findings: Five of nine participants met the eligibility criteria and submitted the survey. All respondents had received care for urgent health needs (5/5) or chronic conditions (4/5). Satisfaction with routine care was found to be high, although trust in providers to respect gender identity and sexual orientation was moderate. The most reported barrier was universal assumption-making by healthcare staff about patients' gender identity or sexual orientation. Securing access to a provider knowledgeable in 2SLGBTQIA+ health needs was also reported to be difficult; however, a trusted provider relationship was the strongest facilitator of identity disclosure.

Discussion: These findings are consistent with patterns documented in 2SLGBTQIA+ health literature, showing a complex intersectionality in sexual orientation, gender orientation, socio-economic status, ethnicity, and other personal/cultural aspects to access care. Barriers such as heteronormative assumptions and a lack of inclusive intake processes continue to block equitable access.

Conclusions: The results suggest gaps in affirming care in North Battleford. Provider education, inclusive intake forms, and investment in primary care continuity are priority areas for improving 2SLGBTQIA+ healthcare equity in this region.

Recommendations: Enhanced education, clinical practices, and institutional policies, along with continual feedback from local PRIDE networks could help ensure the delivery of equitable and inclusive healthcare in the community.

References:

1. Bell J, Purkey E. Trans Individuals' Experiences in Primary Care. *Can Fam Physician*. 2019 Apr;65(4):e147-154.
2. Christopherson L, McLaren K, Schramm L, Holinaty C, Ruddy G, Boughner E, et al. Assessment of Knowledge, Comfort, and Skills Working with Transgender Clients of Saskatchewan Family Physicians, Family Medicine Residents, and Nurse Practitioners. *Transgender Health*. 2022 Oct;7(5):468-472.
3. Comeau D, Johnson C, Bouhamdani N. Review of Current 2SLGBTQIA+ Inequities in the Canadian Health Care System. *Front Public Health* [Internet]. 2023 Jul 18 [cited 2026 May 14];11:[13 p.]. Available from: <https://doi.org/10.3389/fpubh.2023.1183284>.
4. Rainbow Health Ontario. Health in Focus: 2SLGBTQ Sexual Health [Internet]. Toronto (ON): Rainbow Health Ontario; [cited 2026 May 14]. 17 p. Available from: <https://www.rainbowhealthontario.ca/wp-content/uploads/2022/03/Health-in-Focus-2SLGBTQ-Sexual-Health.pdf>.
5. Rai Y, Yogaparan T. Specific Health Needs for LGBTQ Seniors [Internet]. Toronto (ON): University of Toronto; 2018 [cited 2026 May 14]. 13 p. Available from: <https://thehub.utoronto.ca/geriatrics/wp-content/uploads/2021/07/Geriatrics-HUB-LGBTQ-seniors-matters-the-most3.pdf>.
6. Rose G, McCarron M, Reid M, Fayant-McLeod T, Gulka E, Young J, et al. Using Focus Groups to Inform a Peer Health Navigator Service for People Who Are Transgender and Gender Diverse in Saskatchewan, Canada. *Health Expect* [Internet]. 2024 Apr 1 [cited 2026 May 14];27(2):[10 p.]. Available from: <https://doi.org/10.1111/hex.14022>.
7. Schwab A, Peter N, Lawson K, Jessani A. “Expectation is Always that the Practitioner Might Not Be Okay with Queerness”: Experiences of LGBTQ+ with the Healthcare System in Saskatchewan, Canada. *J Homosex*. 2024 Jan 2;71(1):56-71.
8. Sullivan EA, Stryker SD, Blaszczyk J, Spielvogel R, Ho T, Kiraly B, et al. Transgender Care Is Family Medicine: A Call to Action. *Fam Med*. 2024 Apr;56(4):229-233.
9. Li K, Frumkin A, Bi WG, Magrill J, Newton C. Biopsy of Canada's Family Physician Shortage. *Fam Med Community Health* [Internet]. 2023 May 12 [cited 2026 May 14];11(2):[4 p.]. Available from: <https://doi.org/10.1136/fmch-2023-002236>.
10. Renner J, Blaszczyk W, Tauber L, Dekker A, Briken P, Nieder TO. Barriers to Accessing Health Care in Rural Regions by Transgender, Non-Binary, and Gender Diverse People: A Case-Based Scoping Review. *Front Endocrinol* [Internet]. 2021 Nov 18 [cited 2026 May 14];12:[12 p.]. Available from: <https://doi.org/10.3389/fendo.2021.717821>.

Investigating parenthood considerations and subsequent challenges while training in a Family Medicine Residency in Saskatchewan

Anum Ali, FMR II; Aliaa El Tobgy, FMR II
Eriny Tawedrous, MBBCh, CCFP; Rhonda Bryce, MD, MSc

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Residency is a demanding phase of training often coinciding with the period when physicians consider starting or expanding their families. In family medicine, the emphasis on continuity and holistic care may conflict with the reality of long hours, emotional demands, and rigid schedules within training. Parenthood decisions in residency are likely influenced by multiple factors including workload, financial pressures, culture of the program, and career progression. Different programs may vary in their parental leave policies, clarity, consistency, and cultural acceptance regarding having a child during residency.

Question(s): This study aimed to explore parenthood considerations, challenges, and perceived program supports among family medicine residents at the University of Saskatchewan, including factors influencing decisions to have children during residency.

Methods/Methodology: This study was approved by the University of Saskatchewan Behavioural Research Ethics Board (REB #5653). An anonymous, online questionnaire was distributed to family medicine and FM+1 residents over one year. The survey included approximately 50 structured questions assessing decision-making factors, experiences of parenthood during residency, and perceptions of program support.

Results/Findings: Forty-one submitted responses were analyzed. Respondents were most often aged 30–34 years (36.6%), identified as cis-gender women (65.9%), and were in partnered relationships (87.8%). Overall, 24.4% reported having children during residency. Experiences among resident parents were generally positive, particularly regarding program leadership and return-to-training support, although leave duration varied, and some desired longer leave. Reported challenges included guilt related to family separation, on-call demands, and loss of peer continuity. Among residents without children, key barriers included timing within training, perceived impact on career progression, and limited childcare access.

Discussion: Findings highlight a complex interplay between training demands and family planning. While supportive elements exist, variability in leave practices, reintegration experiences, continuing medical education, and childcare access suggests inconsistency in policy implementation.

Conclusions: Residents at the University of Saskatchewan's family medicine program experience both meaningful support and ongoing structural challenges when navigating parenthood during training.

Recommendations: Parental leave policies should promote consistent and equitable support for residents across sites, while allowing implementation flexibility. Enhancing scheduling flexibility, expanding childcare access, and strengthening reintegration supports may help address current gaps and improve training environments.

References:

1. Sullivan MC, Yeo H, Roman SA, Bell RH Jr, Sosa JA. Striving for Work-Life Balance: Effect of Marriage and Children on the Experience of 4402 US General Surgery Residents. *Ann Surg.* 2013 Mar;257(3):571-576.
2. Stack SW, Eurich KE, Kaplan EA, Ball AL, Mookherjee S, Best JA. Parenthood During Graduate Medical Education: A Scoping Review. *Acad Med.* 2019 Nov;94(11):1814-1824.
3. Lavergne MR, Gonzalez A, Ahuja MA, Hedden L, McCracken R. The Relationship Between Gender, Parenthood and Practice Intentions Among Family Medicine Residents: Cross-Sectional Analysis of National Canadian Survey Data. *Hum Resour Health [Internet].* 2019 Aug 15 [cited 2026 May 13];17(1):[16 p.]. Available from: <https://doi.org/10.1186/s1260-19-0396-y>.
4. Morris L, Lindbloom E, Kruse R, Washington KT, Cronk NJ, Paladine HL. Perceptions of Parenting Residents Among Family Medicine Residency Directors. *Fam Med.* 2018 Nov;50(10):756-762.
5. Jakubowski JS, Baltzer H, Lipa JE, Snell L. Parental-Leave Policies and Perceptions of Pregnancy During Surgical Residency Training in North America: A Scoping Review. *Can J Surg.* 2023 Mar;66(2):E132-138.
6. Dason ES, Kapsack A, Baxter NN, Gesink D, Shapiro H, Simpson AN. Resident and Fellow Perspectives on Family Planning and Building During Training. *JAMA Netw Open [Internet].* 2024 Aug 1 [cited 2026 May 13];7(8):[11 p.]. Available from: <https://doi.org/10.1001/jamanetworkopen.2024.29345>.
7. Stratton T, Cook-Chaimowitz L, Pardhan A, Snelgrove N, Chan TM. Parental Leave Policies in Canadian Residency Education. *J Grad Med Educ.* 2021 Apr;13(2):206-212.
8. Willoughby K, Julien M, Zendel BR, Curran V. National Survey of Canadian Residents and Program Directors Regarding Parental Leave During Residency. *Can Med Educ J.* 2020 Sep;11(5):e16-e30.
9. Walsh A, Gold M, Jensen P, Jedrzkiewicz M. Motherhood During Residency Training: Challenges and Strategies. *Can Fam Physician.* 2005 Jul;51(7):990-991.
10. Stack SW, McKinney CM, Spiekerman C, Best JA. Childbearing and Maternity Leave in Residency: Determinants and Well-Being Outcomes. *Postgraduate Med J.* 2018 Dec;94(1118):694-699.

Influence of prenatal hand expression on postpartum breastfeeding

Dilanjani Pietersz, FMR II; Onyinyechi Nwoke, FMR II;
Anneme Dunhin, MBChB, CCFP; Rhonda Bryce, MD, MSc

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Prenatal hand-expression of colostrum has been promoted in the Battlefords since 2016 through distribution of collection kits at 36 weeks' gestation, but its impact on breastfeeding outcomes remains unassessed. Early breastfeeding is critical, especially in rural settings with access and equity challenges.

Question(s): This study aimed to evaluate whether prenatal hand-expression is associated with (1) increased initiation of breastfeeding within 24 hours postpartum, (2) improved maternal breastfeeding confidence, and (3) differences in obstetrical outcomes. A secondary objective was to assess uptake of hand-expression and its relationship with access to colostrum kits, addressing effectiveness, access, and equity.

Methods/Methodology: A retrospective cohort study was conducted among postpartum patients at Battlefords Union Hospital. Participants completed a survey either electronically or verbally during their hospital stay. Data were collected over 6 months and analyzed using descriptive and comparative statistics. Ethics approval was obtained from the University of Saskatchewan's Behavioral Research Ethics Board (Beh ID 5472).

Results/Findings: Of 77 patients approached, 71 participated; 31 (43.7%) had practiced prenatal hand-expression. Exclusive breastfeeding at time of survey was significantly higher among those who hand-expressed compared to those who did not (64.5% vs. 32.5%, $p=0.007$). High breastfeeding confidence was reported by 93.5% of those who hand-expressed versus 80.0% among those who did not ($p=0.17$), with 77% of those expressing reporting improved confidence. Participants who received a colostrum kit were significantly more likely to practice hand-expression compared to non-recipients (61% vs. 5%, $p<0.001$). However, 22 participants (31.0%) did not receive a kit, associated with lower education and later prenatal care. No significant association was observed between prenatal hand-expression and obstetrical outcomes.

Discussion: Prenatal hand-expression is associated with improved early breastfeeding continuance and may enhance maternal confidence. Kit distribution appears to be the primary driver of uptake, with accessibility linked to social inequities.

Conclusions: Prenatal hand-expression represents an effective, low-risk, and scalable intervention to improve early breastfeeding outcomes in a rural population.

Recommendations: All eligible patients should be offered a colostrum kit, with attention to equitable distribution. Additional support should be provided for those unsuccessful with hand-expression, and strategies to improve access for underserved populations are needed.

References:

1. Lyons KE, Ryan CA, Dempsey EM, Ross RP, Stanton C. Breast Milk, A Source of Beneficial Microbes and Associated Benefits for Infant Health. *Nutrients* [Internet]. 2020 Apr 9 [cited 2026 May 11];12(4):[30 p.]. Available from: <https://doi.org/10.3390/nu12041039>.
2. Forster DA, Moorhead AM, Jacobs SE, Davis PG, Walker SP, McEgan KM, et al. Advising Women with Diabetes in Pregnancy to Express Breastmilk in Late Pregnancy (DAME): A Multicentre, Unblinded, Randomised Controlled Trial. *Lancet*. 2017 Jun;389(10085):2204-2213.
3. Zaman F, Morgan S, Scalora C, Nelson M, Francis J. Prenatal Breastfeeding Education with or without Hand Expressing Human Milk and Breastfeeding Duration in a Rural Population. *Nutrients* [Internet]. 2024 Sep 29 [cited 2026 May 11];16(19):[8 p.]. Available from: <https://doi.org/10.3390/nu16193303>.
4. Mendu SB, Neela AR, Tammali S, Kotha R. Impact of Early Bonding During the Maternal Sensitive Period on Long-Term Effects: A Systematic Review. *Cureus* [Internet]. 2024 Jan 31 [cited 2026 May 11];16(2):[7 p.]. Available from: <https://doi.org/10.7759/cureus.53318>.
5. Victora CG, Bahl R, Barros AJD, França GVA, Horton S, Krasevec J, et al. Breastfeeding in the 21st Century: Epidemiology, Mechanisms, and Lifelong Effect. *Lancet*. 2016 Jan;387(10017):475-490.
6. McCoy MB, Heggie P. In-Hospital Formula Feeding and Breastfeeding Duration. *Pediatrics* [Internet]. 2020 Jul 1 [cited 2026 May 11];146(1):[11 p.]. Available from: <https://doi.org/10.1542/peds.2019-2946>.
7. Chantry CJ, Dewey KG, Peerson JM, Wagner EA, Nommsen-Rivers LA. In-Hospital Formula Use Increases Early Breastfeeding Cessation Among First-Time Mothers Intending to Exclusively Breastfeed. *J Pediatr*. 2014 Jun;164(6):1339-45.
8. Dennis CL. Theoretical Underpinnings of Breastfeeding Confidence: A Self-Efficacy Framework. *J Hum Lact*. 1999 Sep;15(3):195-201.
9. McFadden A, Gavine A, Renfrew MJ, Wade A, Buchanan P, Taylor JL, et al. Support for Healthy Breastfeeding Mothers with Healthy Term Babies. *Cochrane Database Syst Rev* [Internet]. 2017 Feb 28 [cited 2026 May 11];2(2):[239 p.]. Available from: <https://doi.org/10.1002/14651858.cd001141.pub5>.
10. Rollins NC, Bhandari N, Hajeebhoy N, Horton S, Lutter CK, Martines JC, et al. Why Invest, and What It Will Take to Improve Breastfeeding Practices? *Lancet*. 2016 Jan;387(10017):491-504.
11. O'Sullivan TA, Ihlein CN. A Brief History of Antenatal Colostrum Expression, and Where to from Here. *Proc Nutr Soc*. 2025 May;84(2):188-195.

Evaluation of a Prenatal and Postpartum Educational Resource at West Winds Primary Health Care Centre

Abeer Nfaileh, FMR II; Tracey Field, MD, CCFP; Katrina Sawchuk, MEd, PhD

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Patients receiving prenatal care in primary care settings frequently have questions regarding investigations, pregnancy milestones, and postpartum care. At West Winds Primary Health Care Centre (WWPHC), the absence of standardized educational materials contributed to inconsistent information delivery and increased non-urgent patient contact. This represents a local quality improvement problem affecting efficiency, patient-centeredness, and provider workload.

Question(s): What are healthcare provider perceptions of a standardized prenatal and postpartum educational resource, and how does its implementation impact clinic workflow, patient preparedness, and communication burden? This project addresses the dimensions of efficiency and patient-centeredness.

Methods/Methodology: This qualitative study used a Plan-Do-Study-Act cycle (PDSA). A pre-survey was used to assess physician usage of maternal educational tools and advice for the development of a standard resource. The resource was then implemented, and a post survey given to iterate the PDSA cycle. Forty-nine surveys were sent to people involved in prenatal care. The educational resources were developed and provided to maternity patients. Survey items assessed perceptions of workflow efficiency, patient preparedness, and non-urgent patient communication. Descriptive statistics were used to summarize responses. An Ethical Exemption was received (Beh E794).

Results/Findings: Eighteen health care practitioners completed the pre-implementation survey and six completed the post-implementation survey. Prior to implementation, 61% of respondents were unsure how patients accessed prenatal educational materials, indicating lack of a standardized approach. Post-implementation, participants reported perceived benefits including reduced non-urgent patient calls and emails, improved patient preparedness, and enhanced planning of prenatal investigations.

Discussion: These findings suggest that a standardized, accessible educational resources may improve clinic efficiency and patient understanding. Existing literature supports the role of prenatal and postpartum education in improving maternal knowledge, self-efficacy, and healthcare engagement. Limitations include small post-implementation sample size.

Conclusions: A QR code-accessible prenatal and printed postpartum educational resource was perceived to improve patient education and clinic workflow at WWPHC.

Recommendations: Future initiatives should include patient feedback, increase clinician uptake, and measure objective outcomes such as communication frequency and visit efficiency. Additional studies evaluating maternal mental health outcomes and postpartum support interventions may further strengthen educational resource development.

References:

1. de los Reyes S, Al-Khaled D, Brito K, Kholmatov M, Farley B, Kuriloff M, et al. AWHONN Postpartum Discharge Education: Patient Knowledge and Sustainability. *BMJ Open Qual* [Internet]. 2022 Nov 4 [cited 2026 May 11];11(4):[7 p.]. Available from: <https://doi.org/10.1136/bmjog-2021-001528>.
2. Hoffman J, Burke L, Kay C, Hlavaty R, Thompson-Wood J, O'Dell D. Standardizing Postpartum Discharge Instructions with an Educational Video: A Quality Improvement Project. *iProceedings* [Internet]. 2019 Oct 2 [cited 2026 May 11];5(1):[2 p.]. Available from: <https://doi.org/10.2196/15245>.
3. Langley GL, Moen R, Nolan KM, Nolan TW, Norman CL, Provost LP. *The Improvement Guide: A Practical Approach to Enhancing Organizational Performance*. 2nd ed. San Francisco (CA): Jossey-Bass; 2009. 512 p.
4. Institute of Medicine Committee on Quality of Health Care in America. *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington (DC): National Academies Press; 2001. 360 p.
5. Sword W, Watt S, Krueger P. Postpartum Health, Service Needs, and Access to Care Experiences of Immigrant and Canadian-Born Women. *J Obstet Gynecol Neonatal Nurs*. 2006 Nov-Dec;35(6):717-727.
6. Yelland J, Sutherland G, Brown SJ. Postpartum Anxiety, Depression and Social Health: Findings from a Population-Based Survey of Australian Women. *BMC Public Health* [Internet]. 2010 Dec 20 [cited 2026 May 11];10:[11 p.]. Available from: <https://doi.org/10.1186/1471-2458-10-771>.
7. Institute for Healthcare Improvement [Internet]. Boston (MA): IHI; c2026. *Model for Improvement*; [cited 2026 May 5];[about 6 screens]. Available from: <https://www.ihl.org/library/model-for-improvement>.
8. O'Connor E, Senger CA, Henninger M, Coppola E, Gaynes BN. Interventions to Prevent Perinatal Depression: Evidence Report and Systematic Review for the US Preventive Services Task Force. *JAMA*. 2019 Feb;321(6):588-601.

Increasing Rates of Cervical Cancer Screening at a Rural Family Medicine Clinic

Oseihioria Alex-Ohunyon, FMR II; Sonya Kim, FMR II
Emmett Harrison, MD, CCFP (EM); Adam Clay, MSc; Brenda Andreas, BSW, RSW

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Cervical cancer is highly preventable through routine screening. Despite this, cervical cancer screening rates remain suboptimal in many rural primary care settings. Barriers identified include limited opportunistic screening, workflow constraints, provider discomfort, and patient access challenges. At the Associate Family Physicians Clinic (AFPC) in Swift Current, the screening rate was 30.1% among eligible patients which is substantially lower than national benchmarks, highlighting a gap in preventive care delivery.

Question(s): To increase cervical cancer screening uptake at AFPC from 30.1% to 68% over a 9-month period improving effectiveness, access, and equity in preventive care delivery.

Methods/Methodology: This Quality Improvement (QI) initiative used the Model for Improvement. A multidisciplinary stakeholder group, including physicians, residents, nursing staff, medical office assistants, and a patient partner, identified barriers and implemented interventions including reminders in the electronic medical record (EMR), staff audit and feedback, opportunistic screening workflows, patient education materials, and expanded nursing involvement in screening. Monthly data was extracted from Accuro EMR and displayed on run charts time. This project was deemed exempt from ethical review by the University of Saskatchewan Biomedical Research Ethics Board (E-Bio-082).

Results/Findings: The baseline cervical cancer screening rate was 30.1%, among approximately 6,000 eligible patients. Following 9 months of interventions, screening rates increased to approximately 34%, representing a 4% improvement. The most effective intervention was Medical Office Assistant (MOA)/Nursing led patient recall and nursing led pap smear clinics.

Discussion: Team-based interventions, including proactive patient outreach and expanded roles for MOAs and nursing staff, were effective in increasing cervical cancer screening rates at AFPC. In particular, expanding the nursing role to perform pap smear offers significant benefits and contributed to improved screening uptake.

Despite these improvements, the overall increase fell short of the target. There is a need to address ongoing barriers and for sustained, scalable strategies to reach under-screened populations.

Conclusions: This QI initiative fell short of goal of 68% cervical cancer screening uptake sustained incremental gains over time, supporting the effectiveness of structured, team-based interventions.

Recommendations: Future improvement efforts may include expanded nursing-led screening, enhanced patient outreach strategies, and evaluation of self-sampling options as system resources allow.

References:

1. Canadian Cancer Society. Canadian Cancer Statistics 2025 [Internet]. Toronto (ON): Canadian Cancer Society; 2025 [cited 2026 May 14]. 111 p. Available from: <https://cancer.ca/en/cancer-information/resources/publications/canadian-cancer-statistics-2025>.
2. World Health Organization [Internet]. Geneva (CE): World Health Organization; c2026. Cervical Cancer Fact Sheet; 2025 Dec 2 [cited 2026 May 11];[about 5 p.]. Available from: <https://www.who.int/news-room/fact-sheets/detail/cervical-cancer>.
3. Yin C, Li XB. Unlocking Early Detection: How Screening Can Save Lives from Cervical Cancer. *World J Clin Oncol* [Internet]. 2025 May 24 [cited 2026 May 14];16(5):[5 p.]. Available from: <https://doi.org/10.5306/wjco.v16.i5.102456>.
4. Fuzzell LN, Perkins RB, Christy SM, Lake PW, Vadaparampil ST. Cervical Cancer Screening in the United States: Challenges and Potential Solutions for Underscreened Groups. *Prev Med (Baltim)* [Internet]. 2021 Mar [cited 2026 May 14];1144:[8 p.]. Available from: <https://doi.org/10.1016/j.yjmed.2020.106400>.
5. Luckett KL. Implementing a Standardized Protocol to Improve Cervical Cancer Screening Rates in Primary Care. *J Am Assoc Nurse Pract*. 2022 Sep;34(9):1077–1082.
6. Popalis ML, Ramirez SI, Leach KM, Granzow ME, Stoltzfus KC, Moss JL. Improving Cervical Cancer Screening Rates: A Scoping Review of Resources and Interventions. *Cancer Causes Control*. 2022 Nov;33(11):1325–1333.
7. Musa J, Achenbach CJ, O’Dwyer LC, Evans CT, McHugh M, Hou L, et al. Effect of Cervical Cancer Education and Provider Recommendation for Screening on Screening Rates: A Systematic Review and Meta-Analysis. *PLoS One* [Internet]. 2017 Sep 5 [cited 2026 May 14];12(9):[28 p.]. Available from: <https://doi.org/10.1371/journal.pone.0183924>.
8. Lurie N, Margolis KL, McGovern PG, Mink PJ, Slater JS. Why Do Patients of Female Physicians Have Higher Rates of Breast and Cervical Cancer Screening? *J Gen Intern Med*. 1997 Jan;12(1):34–43.
9. King E, Busolo D. The Role of Primary Care Nurse Practitioners in Reducing Barriers to Cervical Cancer Screening: A Literature Review. *Can Oncol Nurs J*. 2022 Apr;32(2):233–244.
10. Paskett ED, Tatum CM, D’Agostino R, Rushing J, Velez R, Michielutte R, et al. Community-Based Interventions to Improve Breast and Cervical Cancer Screening: Results

of the Forsyth County Cancer Screening (FoCaS) Project. *Cancer Epidemiol Biomarkers Prev.* 1999 May;8(5):453–459.

11. Kohnen JB. *The Improvement Guide: A Practical Approach to Enhancing Organizational Performance* [book review]. *Qual Manag J.* 1997 Jan;4(4):85–86.

The Effect of Non-Sterile Gloves on Infection Rates in Family Medicine Office-Based Procedures

Ololade Akinlabi, FMR II; Sarina Sadeghi, FMR II
Andrea Vasquez Carmargo, MD, MSc, CCFP; Adam Clay, MSc
Clara Rocha Michaels, MD, CCFP; Asma Gargoum, BA, MBChB, CCFP

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: The use of sterile gloves for minor procedures in primary care is common practice; however, evidence supporting their superiority over non-sterile gloves remains limited. Several studies suggest comparable infection rates, but data from academic family medicine settings are scarce. This study evaluates whether non-sterile gloves are associated with increased post-procedural infection rates in a primary care office setting.

Question(s): Does the use of non-sterile gloves for in-office procedures result in any difference in infection rate when compared with procedures done with sterile gloves?

Methods/Methodology: We performed a retrospective chart review of patients who had procedures done at the Family Medicine Unit from 2015- 2021. Patient demographics (age, sex, comorbidities), lesion sites, and glove type were extracted. Our primary outcome was the presence of signs of infection at follow-up, or the need for antibiotics at subsequent visits for the purpose of treating surgical wound infections. The infection rates were then calculated. Two-sided 95% confidence intervals for the difference in infection rate were used to assess non-inferiority. To account for patients lost to follow up, an additional sensitivity analysis was performed. We obtained ethical approval through the Saskatchewan Health Authority's Research Ethics Board (SHA-REB-19-85) and also received SHA Operational Approval (OA-SHA-19-85).

Results/Findings: A total of 257 patients (158 non-sterile, 99 sterile) and 311 lesions were included. Among lesions with follow-up, infection rates were 8.8% (10/113) in the non-sterile group and 12.2% (10/82) in the sterile group (difference -3.3%, 95% CI -13.0% to 5.3%, $p=0.447$). Sensitivity analyses accounting for patients lost to follow-up demonstrated similar findings, with consistently small, non-significant differences between groups.

Discussion: The observed differences in infection rates were small and not statistically significant and remained within a clinically acceptable margin. These findings are consistent with prior literature suggesting that non-sterile gloves may provide comparable infection rate in minor procedures.

Conclusions: In this academic family medicine office-based procedure setting, the use of non-sterile gloves was not associated with a higher infection rate compared with sterile gloves.

References:

1. Steve E, Lindblad AJ, Allan GM. Non-Sterile Gloves in Minor Lacerations and Excisions? *Can Fam Physician*. 2017 Mar;63(3):217.
2. Heal C, Sriharan S, Buttner PG, Kimber D. Comparing Non-Sterile to Sterile Gloves for Minor Surgery: A Prospective Randomised Controlled Non-Inferiority Trial. *Med J Aust*. 2015 Jan;202(1):27-31.
3. Rogues AM, Lasheras A, Amici JM, Guillot P, Beylot C, Taïeb A, Gachie JP. Infection Control Practices and Infectious Complications in Dermatological Surgery. *J Hosp Infect*. 2007 Mar;65(3):258-263.
4. Shrestha O, Basukala S, Bhugai N, Bohara S, Bhatt A, Thapa N, et al. Postprocedural Infection Rate After Minor Surgical Procedures Performed with and without Sterile Gloves: A Systematic Review and Meta-Analysis. *Int J Surg*. 2024 Nov;110(11):7341-7352.
5. Dekker M, van Mansfeld R, Borgert M, Maaskant J, Paulus F, Seller A, Jongerden IP. The Combined Application of Hand Hygiene and Non-Sterile Gloves by Nurses in a Tertiary Hospital: A Multi Methods Study. *Antimicrob Resist Infect Control* [Internet]. 2024 Feb 28 [cited 2026 May 14];13(1):[8 p.]. Available from: <https://doi.org/10.1186/s13756-024-01378-5>.
6. Fukushima M, Imamura M, Ito M, Muraoka K, Fukasawa M, Kumagai M, et al. Sterile Versus Non-Sterile Gloves During Cystoscopy: A Randomized Prospective Single-Blind Study. *BJUI Compass*. 2024 Sep;5(1):29-33.
7. Vidya B, Deshmukh S, Sharma SD, Harshini G, Zeeshan M. Microbiological Quality of Surgical and Clean Examination Gloves in Oral Surgery. *Eur J Mol Clin Med*. 2020 Jul;7(9):1210-1218.
8. Friedlander MP, Castelli G. Comparing Rates of Infections Between Sterile and Nonsterile Gloves for Common Outpatient Procedures. *Evid Based Pract*. 2018 Jul;21(7):6.
9. Rietz A, Barzin A, Jones K, Mounsey A. Sterile or Non-Sterile Gloves for Minor Skin Excisions? *J Fam Pract*. 2015 Nov;64(11):723-727.
10. Wang D, Shinder R. Comparison of Postoperative Infection Rates After Office-based Oculoplastic Procedures Using Sterile and Clean Gloves. *Ophthalmic Plast Reconstr Surg*. 2023 Mar-Apr;39(2):146-149.
11. Ho C, Jones A. Disposable, Non-Sterile Gloves for Minor Surgical Procedures: A Review of Clinical Evidence. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health; 2017. 10 p.
12. Michener M, Xia Y, Larrimore D, McGraw T, McCarthy S. A Comparison of Infection Rates During Skin Cancer Excisions Using Nonsterile vs Sterile Gloves: A Prospective Randomized Pilot Study. *J Cosmet Dermatol*. 2019 Oct;18(5):1475-1478.

Artificial Intelligence in Primary Care: A Scoping Review of Perceptions, Adoption, and Emerging Implications of Generative AI

Carina Chan, FMR II; Aathmika Kirubaharan, FMR II;
Jared Oberkirsch, MD, CCFP; Udoka Okpalauwaekwe, MBBS, MPH, PhD

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Artificial intelligence (AI) is increasingly becoming integrated into primary care, however, its adoption is shaped by clinician and patient perceptions, workflow, and the evolving ethical and system-level considerations. Also, the rapid emergence of generative AI have introduced additional complexities that remain poorly understood within primary care contexts.

Question(s): This scoping review examines the perceptions, barriers, and facilitators influencing AI adoption in primary care among clinicians and patients, with attention to the ethical, relational, and system-level factors shaping these experiences.

Methods/Methodology: We conducted a Scoping Review following PRISMA-ScR guidelines. Six databases (MEDLINE, PubMed, Embase, Scopus, Web of Science, and CINAHL) were searched for studies published from 2020 onward, supplemented by grey literature and targeted hand-searching. Eligible sources included empirical studies, reviews, and selected commentaries examining AI in primary care. Data were synthesized using descriptive and thematic analysis. Ethics approval was not applicable for this study.

Results/Findings: Thirty-two studies were included, most from North America and employing qualitative or mixed-methods designs. Clinician perceptions were mixed, with reported benefits including improved efficiency, documentation support, and clinical decision-making. Concerns of AI use centred on trust, accuracy, workflow integration, and medico-legal accountability. Patient perspectives were generally cautious but supportive when clinician oversight was maintained. Generative AI, particularly ambient documentation systems, demonstrated workflow benefits but introduced risks such as hallucinated outputs, authorship ambiguity, and increased verification burden. Ethical, relational, and system-level challenges, including governance, infrastructure, and equity, emerged as consistent cross-cutting themes.

Discussion: AI adoption in primary care is conditional, shaped by trust, workflow fit, and the preservation of clinician–patient relationships. Generative AI represents a notable shift, contributing directly to clinical documentation while introducing new challenges related to accuracy, accountability, and the human dimensions of care. Equity in AI use remains underexplored, with limited representation of diverse patient populations.

Conclusions: AI holds dual potential in primary care, to improve efficiency and reduce clinician burden, while also risking disruption to the relational foundations of care.

Recommendations: Future research should prioritize longitudinal, outcome-based, and equity-focused studies to better understand how AI is experienced across diverse populations and its long-term impact on clinical practice.

References:

1. Blease C, Bernstein MH, Gaab J, Kaptchuk TJ, Kossowsky J, Mandl KD, et al. Computerization and the Future of Primary Care: A Survey of General Practitioners in the UK. PLoS One [Internet]. 2018 Dec 12 [cited 2026 May 13];13(12):[12 p.]. Available from: <https://doi.org/10.1371/journal.pone.0207418>.
2. Angus DC, Khera R, Lieu T, Liu V, Ahmad FS, Anderson B, et al. AI, Health, and Health Care Today and Tomorrow: The JAMA Summit Report on Artificial Intelligence. JAMA. 2025 Nov;334(18):1650-1664.
3. Kelly CJ, Karthikesalingam A, Suleyman M, Corrado G, King D. Key Challenges for Delivering Clinical Impact with Artificial Intelligence. BMC Med [Internet]. 2019 Oct 29 [cited 2026 May 13];17(1):[9 p.]. Available from: <https://doi.org/10.1186/s12916-019-1426-2>.
4. Topol EJ. High-Performance Medicine: The Convergence of Human and Artificial Intelligence. Nat Med. 2019 Jan;25(1):44-56.
5. Terry AL, Kueper JK, Beleno R, Brown JB, Cejic S, Dang J, et al. Is Primary Health Care Ready for Artificial Intelligence? What Do Primary Health Care Stakeholders Say? BMC Med Inform Decis Mak [Internet]. 2022 Sep 9 [cited 2026 May 13];22(1):[11 p.]. Available from: <https://doi.org/10.1186/s12911-022-01984-6>.
6. Bazemore A, Grunert T. Sailing the 7C's: Starfield Revisited as a Foundation of Family Medicine Residency Redesign. Fam Med. 2021 Jul;53(7):506-515.
7. Starfield B, Shi L, Macinko J. Contribution of Primary Care to Health Systems and Health. Milbank Q. 2005 Sep;83(3):457-502.
8. Kocaballi AB, Ijaz K, Laranjo L, Quiroz JC, Rezazadegan D, Tong HL, et al. Envisioning an Artificial Intelligence Documentation Assistant for Future Primary Care Consultations: A Co-Design Study with General Practitioners. J Am Med Inform Assoc. 2020 Nov;27(11):1695-1704.
9. Shah SJ, Crowell T, Jeong Y, Devon-Sand A, Smith M, Yang B, et al. Physician Perspectives on Ambient AI Scribes. JAMA Netw Open [Internet]. 2025 Mar 3 [cited 2026 May 13];8(3):[9 p.]. Available from: <https://doi.org/10.1001/jamanetworkopen.2025.1904>.
10. Shaw J, Agarwal P, Desveaux L, Palma DC, Stamenova V, Jamieson T, et al. Beyond "Implementation": Digital Health Innovation and Service Design. NPJ Digit Med [Internet]. 2018 Sep 20 [cited 2026 May 13];1:[5 p.]. Available from: <https://doi.org/10.1038/s41746-018-0059-8>.
11. Bender EM, Gebu T, McMillan-Major A, Shmitchell S. On the Dangers of Stochastic Parrots: Can Language Models Be Too Big? In: Proceedings of the 2021 ACM Conference on Fairness, Accountability, and Transparency; 2021 Mar 3-10; Virtual. New York (NY): Association for Computing Machinery; c2021. p. 610-623.

12. London AJ. Artificial Intelligence and Black-Box Medical Decisions: Accuracy Versus Explainability. *Hastings Cent Rep*. 2019 Jan;49(1):15-21.
13. Martin J, Rueda A, Lee GH, Tassone VK, Park H, Ivanov M, et al. Digital Interventions to Understand and Mitigate Stress Response: Protocol for Process and Content Evaluation of a Cohort Study. *JMIR Res Protoc* [Internet]. 2024 May 6 [cited 2026 May 13];13:[14 p.]. Available from: <https://doi.org/10.2196/54180>.
14. Kerasidou A, Kerasidou CX. Epistemic Authority and Medical AI: Epistemological Differences and Challenges in Medical Practice. *Med Health Care Philos*. 2026 Mar;29(1):89-95.
15. Babu J, Joseph D, Kumar RM, Alexander E, Sasi R, Joseph J. Emotional AI and the Rise of Pseudo-Intimacy: Are We Trading Authenticity for Algorithmic Affection? *Front Psychol* [Internet]. 2025 Sep 18 [cited 2026 May 13];16:[8 p.]. Available from: <https://doi.org/10.3389/fpsyg.2025.1679324>.
16. Nazer LH, Zatarah R, Waldrip S, Ke JXC, Moukheiber M, Khanna AK, et al. Bias in Artificial Intelligence Algorithms and Recommendations for Mitigation. *PLoS Digit Health* [Internet]. 2023 Jun 22 [cited 2026 May 13];2(6):[14 p.]. Available from: <https://doi.org/10.1371/journal.pdig.0000278>.
17. Levac D, Colquhoun H, O'Brien KK. Scoping Studies: Advancing the Methodology. *Implement Sci* [Internet]. 2010 Sep 20 [cited 2026 May 13];5:[9 p.]. Available from: <https://doi.org/10.1186/1748-5908-5-69>.
18. Peters M, Godfrey C, McInerney P, Munn Z, Tricco AC, Khalil H. Best Practice Guidance and Reporting Items for the Development of Scoping Review Protocols. *JBI Evid Synth*. 2022 Apr;20(4):953-968.
19. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med*. 2018 Oct;169(7):467-473.
20. Thomas J, Harden A. Methods for the Thematic Synthesis of Qualitative Research in Systematic Reviews. *BMC Med Res Methodol* [Internet]. 2008 Jul 10 [2026 May 13];8:[10 p.]. Available from: <https://doi.org/10.1186/1471-2288-8-45>.
21. Alanzi, T., et al., Barriers and Facilitators of Artificial Intelligence in Family Medicine: An Empirical Study With Physicians in Saudi Arabia. *Cureus*, 2023. 15(11): p. e49419.
22. Alanzi T, Alotaibi R, Alajmi R, Bukhamsin Z, Fadaq K, AlGhamdi N, et al. Barriers and Facilitators of Artificial Intelligence in Family Medicine: An Empirical Study with Physicians in Saudi Arabia. *Cureus* [Internet]. 2023 Nov 26 [cited 2026 May 13];15(11):[11 p.]. Available from: <https://doi.org/10.7759/cureus.49419>.
23. Buck C, Doctor E, Hennrich J, Jöhnk J, Eymann T. General Practitioners' Attitudes Toward Artificial Intelligence-Enabled Systems: Interview Study. *J Med Internet Res* [Internet]. 2022 Jan 27 [cited 2026 May 13];24(1):[18 p.]. Available from: <https://doi.org/10.2196/28916>.

24. Catalina QM, Fuster-Casanovas A, Vidal-Alaball J, Escalé-Besa A, Marin-Gomez FX, Femenia J, Solé-Casals J. Knowledge and Perception of Primary Care Healthcare Professionals on the use of Artificial Intelligence as a Healthcare Tool. *Digit Health* [Internet]. 2023 Jun 14 [cited 2026 May 13];9:[11 p.]. Available from: <https://doi.org/10.1177/20552076231180511>.
25. Darcel K, Upshaw T, Craig-Neil A, Macklin J, Steele Gray C, Chan TCY, et al. Implementing Artificial Intelligence in Canadian Primary Care: Barriers and Strategies Identified Through a National Deliberative Dialogue. *PLoS One* [Internet]. 2023 Feb 27 [cited 2026 May 13];18(2):[14 p.]. Available from: <https://doi.org/10.1371/journal.pone.0281733>.
26. Nash DM, Thorpe C, Brown JB, Kueper JK, Rayner J, Lizotte DJ, et al. Perceptions of Artificial Intelligence Use in Primary Care: A Qualitative Study with Providers and Staff of Ontario Community Health Centres. *J Am Board Fam Med*. 2023 Apr;36(2):221-228.
27. Ruan EL, Alkattan A, Elhadad N, Rossetti SC. Clinician Perceptions of Generative Artificial Intelligence Tools and Clinical Workflows: Potential Uses, Motivations for Adoption, and Sentiments on Impact. *AMIA Annu Symp Proc*. 2025 May;2024:960-969.
28. Upshaw TL, Craig-Neil A, Macklin J, Gray CS, Chan TCY, Gibson J, Pinto AD. Priorities for Artificial Intelligence Applications in Primary Care: A Canadian Deliberative Dialogue with Patients, Providers, and Health System Leaders. *J Am Board Fam Med*. 2023 Apr;36(2):210-220.
29. Waheed MA, Liu L. Perceptions of Family Physicians About Applying AI in Primary Health Care: Case Study from a Premier Health Care Organization. *JMIR AI* [Internet]. 2024 Apr 17 [cited 2026 May 13];3:[17 p.]. Available from: <https://doi.org/10.2196/40781>.
30. Wiedermann CJ, Mahlkecht A, Piccoliori G, Engl A. Redesigning Primary Care: The Emergence of Artificial-Intelligence-Driven Symptom Diagnostic Tools. *J Pers Med* [Internet]. 2023 Sep 15 [cited 2026 May 13];13(9):[7 p.]. Available from: <https://doi.org/10.3390/jpm13091379>
31. Olson KD, Meeker D, Troup M, Barker TD, Nguyen VH, Manders JB, et al. Use of Ambient AI Scribes to Reduce Administrative Burden and Professional Burnout. *JAMA Netw Open* [Internet]. 2025 Oct 1 [cited 2026 May 13];8(10):[12 p.]. Available from: <https://doi.org/10.1001/jamanetworkopen.2025.34976>.
32. Lukac PJ, Turner W, Vangala S, Chin AT, Khalili J, Shih YT, et al. Ambient AI Scribes in Clinical Practice: A Randomized Trial. *NEJM AI* [Internet]. 2025 Nov 26 [cited 2026 May 13];2(12):[18 p.]. Available from: <https://doi.org/10.1056/aioa2501000>
33. van Linschoten RCA, van Loon CM, Joanknecht L, Bischoff EWMA. Ambient Scribe in General Practice: A Multi-Perspective Before-After Longitudinal Mixed-Methods Study. *NPJ Digit Med* [Internet]. 2026 Mar 2 [cited 2026 May 13];9(1):[9 p.]. Available from: <https://doi.org/10.1038/s41746-026-02454-3>.
34. Ho V, Brown Johnson C, Ghanzouri I, Amal S, Asch S, Ross E. Physician- and Patient-Elicited Barriers and Facilitators to Implementation of a Machine Learning-Based

- Screening Tool for Peripheral Arterial Disease: Preimplementation Study with Physician and Patient Stakeholders. *JMIR Cardio* [Internet]. 2023 Nov 6 [cited 2026 May 13];7:[10 p.]. Available from: <https://doi.org/10.2196/44732>
35. Rotenstein L, Wang L, Zupanc SN, Penumarthy A, Laurentiev J, Lamey J, et al. Looking Beyond Mortality Prediction: Primary Care Physician Views of Patients' Palliative Care Needs Predicted by a Machine Learning Tool. *Appl Clin Inform*. 2024 May;15(3):460-468.
 36. Tanguay-Sela M, Benrimoh D, Popescu C, Perez T, Rollins C, Snook E, et al. Evaluating the Perceived Utility of an Artificial Intelligence-Powered Clinical Decision Support System for Depression Treatment Using a Simulation Center. *Psychiatry Res* [Internet]. 2022 Feb 1 [cited 2026 May 13];308:[9 p.]. Available from: <https://doi.org/10.1016/j.psychres.2021.114336>.
 37. Stroud AM, Curtis SH, Weir IB, Stout JJ, Barry BA, Bobo WV, et al. Physician Perspectives on the Potential Benefits and Risks of Applying Artificial Intelligence in Psychiatric Medicine: Qualitative Study. *JMIR Ment Health* [Internet]. 2025 Feb 10 [cited 2026 May 13];12:[11 p.]. Available from: <https://doi.org/10.2196/64414>.
 38. Ghadiri P, Yaffe MJ, Adams AM, Abbasgholizadeh-Rahimi S. Primary Care Physicians' Perceptions of Artificial Intelligence Systems in the Care of Adolescents' Mental Health. *BMC Prim Care* [Internet]. 2024 Jun 13 [cited 2026 May 13];25(1):[14 p.]. Available from: <https://doi.org/10.1186/s12875-024-02417-1>.
 39. Chew HSJ, Achananuparp P. Perceptions and Needs of Artificial Intelligence in Health Care to Increase Adoption: Scoping Review. *J Med Internet Res* [Internet]. 2022 Jan 14 [cited 2026 May 13];24(1):[19 p.]. Available from: <https://doi.org/10.2196/32939>.
 40. Garies S, Liang S, Weyman K, Durant S, Ramji N, Alhaj M, Pinto A. Artificial Intelligence in Primary Care Practice: Qualitative Study to Understand Perspectives on Using AI to Derive Patient Social Data. *Can Fam Physician*. 2024 Jul-Aug;70(7-8):e102-109.
 41. Garies S, Liang S, Weyman K, Ramji N, Alhaj M, Pinto AD. Developing an AI Tool to Derive Social Determinants of Health for Primary Care Patients: Qualitative Findings from a Codesign Workshop. *Ann Fam Med*. 2024 Jul-Aug;22(4):317-324.
 42. Mahlkecht A, Engl A, Piccoliori G, Wiedermann CJ. Supporting Primary Care Through Symptom Checking Artificial Intelligence: A Study of Patient and Physician Attitudes in Italian General Practice. *BMC Prim Care* [Internet]. 2023 Sep 4 [cited 2026 May 13];24(1):[18 p.]. Available from: <https://doi.org/10.1186/s12875-023-02143-0>.
 43. Leiserowitz G, Mansfield J, MacDonald S, Jost M. Patient Attitudes Toward Ambient Voice Technology: Preimplementation Patient Survey in an Academic Medical Center. *JMIR Med Inform* [Internet]. 2025 Nov 27 [cited 2026 May 13];13:[10 p.]. Available from: <https://doi.org/10.2196/77901>.
 44. Tu T, Schaekermann M, Palepu A, Saab K, Freyberg J, Tanno R, et al. Towards Conversational Diagnostic Artificial Intelligence. *Nature*. 2025 Jun;642(8067):442-450.

45. Nong P, Neprash HT. Unintended Consequences of Using Ambient Artificial Intelligence Scribes for Billing. *JAMA Health Forum* [Internet]. 2026 Jan 2 [cited 2026 May 13];7(1):[3 p.]. Available from: <https://doi.org/10.1001/jamahealthforum.2025.5771>.
46. Scipion CEA, Manchester MA, Federman A, Wang Y, Arias JJ. Barriers to and Facilitators of Clinician Acceptance and Use of Artificial Intelligence in Healthcare Settings: A Scoping Review. *BMJ Open* [Internet]. 2025 Apr 15 [cited 2026 May 13];15(4):[13 p.]. Available from: <https://doi.org/10.1136/bmjopen-2024-092624>.
47. Schragger S, Seehusen DA, Sexton SM, Richardson CR, Neher JO, Pimlott N, et al. Use of AI in Family Medicine Publications: A Joint Editorial from Journal Editors. *Fam Med*. 2025 Jan;57(1):1-5.
48. Ayaz A, Yanartaş M. An Analysis on the Unified Theory of Acceptance and Use of Technology Theory (UTAUT): Acceptance of Electronic Document Management System (EDMS). *Comput Hum Behav Rep* [Internet]. 2020 Oct 10 [cited 2026 May 13];2:[7 p.]. Available from: <https://doi.org/10.1016/j.chbr.2020.100032>.
49. Chustecki M. Benefits and Risks of AI in Health Care: Narrative Review. *Interact J Med Res* [Internet]. 2024 Nov 18 [cited 2026 May 13];13:[13 p.]. Available from: <https://doi.org/10.2196/53616>.
50. Damji AN, Martin D, Lermen N Jr, Pinto LF, Trindade TGD, Prado JC Jr. Trust As the Foundation: Thoughts on the Starfield Principles in Canada and Brazil: The Besroul Papers: A Series on the State of Family Medicine in Canada and Brazil. *Can Fam Physician*. 2018 Nov;64(11):811-815.
51. Hedden L, Spencer S, Mathews M, Gard Marshall E, Lukewich J, Asghari S, et al. "Technology Has Allowed Us to Do a Lot More but It's Not Necessarily the Panacea for Everybody": Family Physician Perspectives on Virtual Care During the COVID-19 Pandemic and Beyond. *PLoS One* [Internet]. 2024 Feb 29 [cited 2026 May 13];19(2):[16 p.]. Available from: <https://doi.org/10.1371/journal.pone.0296768>.
52. Blease C, Hagström J, Sanchez CG, Kharko A, McMillan B, Gaab J, et al. General Practitioners' Adoption of Generative Artificial Intelligence in Clinical Practice in the UK: An Updated Online Survey. *Digit Health* [Internet]. 2025 Nov 25 [cited 2026 May 13];11:[9 p.]. Available from: <https://doi.org/10.1177/20552076251394287>
53. Bajwa J, Munir U, Nori A, Williams B. Artificial Intelligence in Healthcare: Transforming the Practice of Medicine. *Future Healthc J*. 2021 Jul;8(2):e188-194.
54. Price WN 2nd, Cohen IG. Privacy in the Age of Medical Big Data. *Nat Med*. 2019 Jan;25(1):37-43.
55. Collier R. Electronic Health Records Contributing to Physician Burnout. *CMAJ*. 2017 Nov;189(45):E1405-1406.
56. Reardon CM, Damschroder LJ, Ashcraft LE, Kerins C, Bachrach RL, Nevedal AL, et al. The Consolidated Framework for Implementation Research (CFIR) User Guide: A Five-Step Guide for Conducting Implementation Research Using the Framework. *Implement Sci*

[Internet]. 2025 Aug 16 [cited 2026 May 13];20(1):[13 p.]. Available from:
<https://doi.org/10.1186/s13012-025-01450-7>.

57. Oblath R, Beaugard CA, Herrera CN, Xu C, Syed S, Sadatis C, et al. Bridging Crisis and Care: Exploring the Role of Behavioral Health Professionals in a Police Co-Response Model. Health Justice [Internet]. 2025 Nov 25 [cited 2026 May 13];13(1):[12 p.]. Available from: <https://doi.org/10.1186/s40352-025-00381-1>.
58. Adler-Milstein J, Aggarwal N, Ahmed M, Castner J, Evans BJ, Gonzalez AA, et al. Meeting the Moment: Addressing Barriers and Facilitating Clinical Adoption of Artificial Intelligence in Medical Diagnosis. NAM Perspect [Internet]. 2022 Sep 29 [cited 2026 May 13];2022:[29 p.]. Available from: <https://doi.org/10.31478/202209c>.

Understanding Missed Appointments at West Winds Primary Health Centre: Perspectives and Opportunities

Stephanie Walls, FMR II; Maa Quartey, FMR II;
Jason Hosain, MD, CCFP; Katrina Sawchuk, MEd, PhD

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Missed appointments represent a persistent challenge in primary care, contributing to inefficiencies, lost capacity, and gaps in patient care. Hitherto, no formal examination of missed appointments has occurred within WWPHC. Understanding the nature, frequency, and contributory factors to missed appointments is important, as optimizing efficiency improves patient experience and outcomes, and staff satisfaction. This project addresses a local quality improvement gap – examining missed appointments at WWPHC and their implications for access, efficiency, and patient-centered care.

Question(s): This study asked: (1) Among WWPHC patients, what patient-level demographic, clinical, and access-related factors are associated with missed family physician appointments? (2) How do these factors differ between patients missing one appointment and those missing multiple appointments? Primarily addressed were quality dimensions of efficiency, access, and patient-centeredness.

Methods/Methodology: A retrospective EMR review examined the most recent 99 missed appointments over a one-month period at WWPHC. Data included demographics, postal codes, comorbidities, psychological factors, and appointment type. Data was analyzed in Excel using descriptive statistics and thematic qualitative analysis. This project received exemption from the University of Saskatchewan Research Ethics Board #E-Bio118.

Results/Findings: Missed appointments studied represented nearly one full working week of family physician time. Most patients missed single appointments; however, some missed multiple. Prescription refills and result follow-ups were most commonly missed. Missed appointments clustered in neighborhoods closest to WWPHC. Fewer missed appointments occurred among patients with household incomes above \$75,000.

Discussion: Findings suggest that commonly missed appointments may represent opportunities for alternative care modalities, particularly virtual care, given their non-acute nature. With the high proportion of unattached patients provincially, addressing inefficiencies associated with missed appointments may improve access and overall system capacity.

Conclusions: Missed appointments at WWPHC are common, result in considerable lost clinical time, and are influenced by appointment type and socioeconomic factors. Addressing this could improve efficiency and support patient-centered care delivery.

Recommendations Future improvement initiatives include implementing EMR-based documentation of missed appointment reasons, expanding reminder and notification supports, testing flexible scheduling and virtual care options for appropriate visits. Examining over-users could provide insight for streamlining patient care for future improvement studies.

References:

1. Parsons J, Bryce C, Atherton H. Which Patients Miss Appointments with General Practice and the Reasons Why: A Systematic Review. *Br J Gen Pract.* 2021 May;71(707):e406-412.
2. Ciechanowski P, Russo J, Katon W, Simon G, Ludman E, et al. Where Is the Patient? The Association of Psychosocial Factors and Missed Primary Care Appointments in Patients with Diabetes. *Gen Hosp Psychiatry.* 2006 Jan-Feb;28(1):9-17.
3. Greig EC, Gonzalez-Colaso R, Nwanyanwu K. Racial, Ethnic, and Socioeconomic Disparities Drive Appointment No-Show in Patients with Chronic Eye Disease. *J Racial Ethn Health Disparities.* 2023 Aug;10(4):1790-1797.
4. Kaplon AW, Lindemann TL, Patel PA, Soliman AMS. Factors Contributing to Late Cancellation and No-Show for Otolaryngologic Surgery: A Prospective Study. *J Healthc Qual.* 2022 Mar-Apr;44(2):88-94.
5. Taheri-Shirazi M, Namdar K, Ling K, Karmali K, McCradden MD, Lee W, Khalvati F. Exploring Potential Barriers in Equitable Access to Pediatric Diagnostic Imaging Using Machine Learning. *Front Public Health* [Internet]. 2023 Feb 24 [cited 2026 May 11];11:[11 p.]. Available from: <https://doi.org/10.3389/fpubh.2023.968319>.
6. McGrath L, Patel K, Garcia IC, Shalen E, Burton Y, Khan A. Factors Associated with Nonattendance at the Adult Congenital Heart Disease Clinic. *Am J Cardiol.* 2023 Aug;201(8):247-251.
7. Samuels RC, Ward VL, Melvin P, Macht-Greenberg M, Wenren LM, Yi J, et al. Missed Appointments: Factors Contributing to High No-Show Rates in an Urban Pediatrics Primary Care Clinic. *Clin Pediatr (Phila).* 2015 Sep;54(10):976-82.
8. Santos J, Camplain C, Pollitt AM, Baldwin JA. A Formative Assessment of Client Characteristics Associated with Missed Appointments in Integrated Primary Care Services in Rural Arizona. *J Eval Clin Pract.* 2024 Mar;30(2):243-250.
9. Adepoju OE, Liaw W, Phillips CD. Longer Appointment Duration Reduces Future Missed Appointments in Safety-Net Clinics. *Am J Manag Care.* 2025 Feb;31(2):e39-46.
10. Teo AR, Niederhausen M, Handley R, Metcalf EE, Call AA, Jacob RL, et al. Using Nudges to Reduce Missed Appointments in Primary Care and Mental Health: A Pragmatic Trial. *J Gen Intern Med.* 2023 Jul;38(Suppl 3):894-904.

Exploring the Relationship Between Clinical Scheduling Practices, Resident Wellness, and Patient Care in Saskatchewan's Family Medicine Training Programs

Samantha Tso, FMR II; Hadir Ismail, FMR I;
Mahmood Beheshti, MD, CCFP; Udoka Okpalauwaekwe, MBBS, MPH, PhD

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Medical residency is characterized by prolonged work hours that contribute to sleep deprivation, fatigue, and burnout, with well-established impacts on cognitive performance, decision-making, and patient care. Although extensive research, largely conducted prior to 80-hour workweek reforms in the United States, has examined these effects, less is known in the Canadian context, where resident work hours are provincially regulated. In Saskatchewan, call structures vary widely across family medicine training sites, with inconsistencies in shift lengths and post-call recovery periods. The purpose of this study was to illuminate residents' perspectives to optimize wellness and learning outcomes.

Question(s): Are higher clinical scheduling demands associated with increased emotional exhaustion among family medicine residents?

Methods/Methodology: This mixed-methods cross-sectional study combined quantitative survey data with open-ended qualitative responses. All PGY-1 and PGY-2 residents at urban, rural, and regional training sites in Saskatchewan's Department of Family Medicine residency program were invited to participate by email. Measures included scheduling burden (weekly work hours, call frequency, post-call recovery), fatigue exposure (frequency of sleep deprivation, sleep during call), emotional exhaustion (adapted from the Maslach Burnout Inventory), and psychological distress (GAD-7). Emotional exhaustion was dichotomized at the sample median. Descriptive statistics and bivariate analyses were conducted, followed by multivariable logistic regression. Ethics approval was obtained from the University of Saskatchewan Research Ethics Board (ID#5885).

Results/Findings: Residents reporting more frequent sleep deprivation and those working more than 60 hours per week were more likely to experience high emotional exhaustion, though these associations were not statistically significant. Psychological distress was strongly associated with emotional exhaustion, and residents reporting moderate-to-severe anxiety were significantly more likely to experience high emotional exhaustion ($p < 0.001$). Call frequency showed an inverse, non-significant association with emotional exhaustion, and other scheduling variables were not significantly associated in bivariate analyses.

Conclusions: Our study's findings suggest a pathway linking scheduling burden, fatigue, and emotional exhaustion. Emotional exhaustion was also linked to perceived impacts on clinical decision-making and patient safety. These findings underscore the importance of addressing structural scheduling practices to support resident well-being, physician retention, and high-quality patient care. Further longitudinal research is needed to inform targeted interventions.

References:

1. Dyrbye LN, Thomas MR, Shanafelt TD. Medical Student Distress: Causes, Consequences, and Proposed Solutions. *Mayo Clin Proc.* 2008 Dec;80(12):1613–1622.
2. Dyrbye LN, West CP, Satele D, Boone S, Tan L, Sloan J, et al. Burnout Among U.S. Medical Students, Residents, and Early Career Physicians Relative to the General U.S. Population. *Acad Med.* 2014 Mar;89(3):443–451.
3. Landrigan CP, Rothschild JM, Cronin JW, Kaushal R, Burdick E, Katz JT. Effect of Reducing Interns' Work Hours on Serious Medical Errors in Intensive Care Units. *New Engl J Med.* 2004 Oct;351(18):1838–1848.
4. Lockley SW, Cronin JW, Evans EE, Cade BE, Lee CJ, Landrigan CP, et al. Effect of Reducing Interns' Weekly Work Hours on Sleep and Attentional Failures. *New Engl J Med.* 2004 Oct;351(18):1829–1837.
5. Maslach C, Leiter MP. Understanding the Burnout Experience: Recent Research and Its Implications for Psychiatry. *World Psychiatry.* 2016 Jun;15(2):103–111.
6. Panagioti M, Geraghty K, Johnson J, Zhou A, Panagopoulou E, Chew-Graham C, et al. Association Between Physician Burnout and Patient Safety, Professionalism, and Patient Satisfaction. *JAMA Intern Med.* 2018 Oct;178(10):1317–1331.
7. Philibert I. Sleep Loss and Performance in Residents and Nonphysicians: A Meta-Analytic Examination. *Sleep.* 2005 Nov;28(11):1392–1402.
8. West CP, Dyrbye LN, Shanafelt TD. Physician Burnout: Contributors, Consequences, and Solutions. *J Intern Med.* 2018 Jun;283(6):516–529.

Exploration of Patient Perspectives of the Use of Artificial Intelligence (AI) Scribe in a Family Medicine Clinic

Riley Schaaf, FMR II; Gezila De Beer, FMR II; Oluwaseun Ajayi, FMR II
Danielle Frost, BSc (Hon.), MD, MEd, CCFP (SEM), Dip. Sport Med (CASEM)
Andries Muller, MBChB, PhD, CCFP, FCFP; Katrina Sawchuk, MEd. PhD

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Ambient artificial intelligence (AI) scribes are increasingly used in primary care to generate draft clinical notes by listening to patient–physician interactions. While they may improve efficiency and workflow, they also raise ethical concerns related to patient comfort, trust, understanding, and consent. Understanding patient perspectives is essential for transparent and ethical implementation.

Question(s): This study explored patients’ perceptions of the use of ambient AI scribes in a primary care clinic, including awareness, comfort, perceived benefits, and concerns. It aimed to collect anonymous perspectives outside the clinical encounter to encourage candid responses.

Methods/Methodology: A mixed-methods design supported voluntary participation and minimized undue influence. Data were collected via a paper survey at clinic reception and an electronic SurveyMonkey® version accessed through a QR code. Participants could opt into a follow-up phone interview with three structured questions, recorded using Zoom®. Data collection occurred over six weeks. Ethics Approval was obtained from the University of Saskatchewan’s Beh # 6026 and SHA OA-UofS-6026.

Results/Findings: Twenty-four participants completed the survey, and four completed interviews. About half (54.2%) had heard of AI scribes, though most reported limited understanding (62.5%). A slight majority (56.5%) were somewhat or very comfortable with their use, and most believed they could improve healthcare quality. Privacy and data security were the most common concerns (75.0%). Perceived benefits included improved physician focus, time savings, and reduced documentation burden. Concerns included trust, accuracy, and reduced human interaction. Many participants expressed a need for greater transparency and education.

Discussion: These findings contribute to emerging research on patient perspectives of AI in healthcare, which has largely focused on clinicians. Generalizability is limited by the single-site setting.

Conclusions: Patients demonstrated cautious acceptance of AI scribes, influenced by limited understanding and concerns about privacy and autonomy. Acceptance was closely tied to being informed and having a choice.

Recommendations: Clinicians should clearly explain AI Scribe use, including data security, accuracy safeguards, and patients’ right to decline, to support trust and informed decision-making.

References:

1. Chandrasekaran R, Moustakas E. Patient Attitudes Toward Ambient Artificial Intelligence Scribes in Clinical Care: Insights from a Cross-Sectional Study. *J Am Med Inform Assoc*. 2026;33(2):263–272.
2. Evans K, Papinniemi A, Ploderer B, Nicholson V, Hindhaugh T, Vuvan V, et al. Impact of Using an AI Scribe on Clinical Documentation and Clinician–Patient Interactions in Allied Health Private Practice: Perspectives of Clinicians and Patients. *Musculoskelet Sci Pract* [Internet]. 2025 Aug 1 [cited 2026 May 12];78:[9 p.]. Available from: <https://doi.org/10.1016/j.msksp.2025.103333>.
3. Sasseville M, Yousefi F, Ouellet S, Naye F, Stefan T, Carnovale V, et al. The Impact of AI Scribes on Streamlining Clinical Documentation: A Systematic Review. *Healthcare (Basel)* [Internet]. 2025 Jun 16 [cited 2026 May 12];13(12):[21 p.]. Available from: <https://doi.org/10.3390/healthcare13121447>.
4. Tierney AA, Gayre G, Hoberman B, Mattern B, Balleca M, Kipnis P, et al. Ambient Artificial Intelligence Scribes to Alleviate the Burden of Clinical Documentation. *NEJM Catal Innov Care Deliv* [Internet]. 2024 Feb 21 [cited 2026 May 12];5(3):[15 p.]. Available from: <https://catalyst.nejm.org/doi/full/10.1056/CAT.23.0404>.
5. Chrysosofos S, Ochoa E, Sacks JM. The Digital Scribe: A New Wave of Efficiency and Quality of Life for Plastic Surgeons. *Plast Reconstr Surg Glob Open* [Internet]. 2025 May 5 [cited 2026 May 12];13(5):[2 p.]. Available from: <https://doi.org/10.1097/gox.00000000000006754>.
6. Moschogianis S, Darley S, Coulson T, Peek N, Cheraghi-Sohi S, Brown BC. Seven Opportunities for Artificial Intelligence in Primary Care Electronic Visits: Qualitative Study of Staff and Patient Views. *Ann Fam Med*. 2025 27;23(3):214–222.
7. Cao DY, Silkey JR, Decker MC, Wanat KA. Artificial Intelligence–Driven Digital Scribes in Clinical Documentation: Pilot Study Assessing the Impact on Dermatologist Workflow and Patient Encounters. *JAAD Int*. 2024 Feb;15(2):149–151.
8. Haberle T, Cleveland C, Snow GL, Barber C, Stookey N, Thornock C, et al. The Impact of Nuance DAX Ambient Listening AI Documentation: A Cohort Study. *J Am Med Inform Assoc*. 2024 Apr;31(4):975–979.
9. Aggarwal R, Farag S, Martin G, Ashrafian H, Darzi A. Patient Perceptions on Data Sharing and Applying Artificial Intelligence to Health Care Data: Cross-Sectional Survey. *J Med Internet Res* [Internet]. 2021 Aug 26 [cited 2026 May 12];23(8):[12 p.]. Available from: <https://doi.org/10.2196/26162>.
10. Foresman G, Biro J, Tran A, MacRae K, Kazi S, Schubel L, et al. Patient Perspectives on Artificial Intelligence in Health Care: Focus Group Study for Diagnostic Communication and Tool Implementation. *J Particip Med* [Internet]. 2025 Jul 24 [cited 2026 May 12];17:[15 p.]. Available from: <https://doi.org/10.2196/69564>.
11. Shah SJ, Crowell T, Jeong Y, Devon-Sand A, Smith M, Yang B, et al. Physician Perspectives on Ambient AI Scribes. *JAMA Netw Open* [Internet]. 2025 Mar 3 [cited 2026 May 12];8(3):[9 p.]. Available from: <https://doi.org/10.1001/jamanetworkopen.2025.1904>.
12. Esmaeilzadeh P, Mirzaei T, Dharanikota S. Patients’ Perceptions Toward Human–Artificial Intelligence Interaction in Health Care: Experimental Study. *J Med Internet*

- Res [Internet]. 2021 Nov 25 [cited 2026 May 12];23(11):[21 p.]. Available from: <https://doi.org/10.2196/25856>.
13. Street RL Jr, Liu L, Farber NJ, Chen Y, Calvitti A, Weibel N, et al. Keystrokes, Mouse Clicks, and Gazing at the Computer: How Physician Interaction with the EHR Affects Patient Participation. *J Gen Intern Med*. 2018 Apr;33(4):423–428.
 14. CBC News [Internet]. Toronto (ON): CBC; c2026. AI-related data centres use vast amounts of water. But gauging how much is a murky business; 2025 Oct 18 [updated 2025 Oct 22; cited 2026 May 10];[about 11 screens]. Available from: <https://www.cbc.ca/news/ai-data-centre-canada-water-use-9.6939684>.
 15. CNN [Internet]. Atlanta (GA): CNN; c2026. Data centers are creating heat islands, warming the land around them by up to 16 degrees; 2026 Mar 30 [cited 2026 May 10];[about 5 screens]. Available from: <https://www.cnn.com/2026/03/30/climate/data-centers-are-having-an-underreported>.

Penicillin Allergy De-labelling at the Family Medicine Unit: A Quality Improvement Project

Kylie Kosokowsky, FMR II; Shaina Templeton, FMR II
Radhika Marwah, MBBS, CCFP, MSc-AMH (Addictions)
Debbie Bunka, BSc (Pharm); Adam Clay, MSc

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: In the general population, 10-15% of people report a penicillin allergy but only 10% of these individuals have a positive penicillin skin test. Delabelling of patients without true allergy can reduce antibiotic resistance and adverse events. The PEN-FAST tool is a point-of-care risk assessment to determine low-risk or high-risk for a true penicillin allergy.

Question(s): The quality improvement project aimed to delabel 10% of adult patients documented at the Family Medicine Unit (FMU) as having a penicillin allergy in 6 months.

Methods/Methodology: The Model for Improvement was used. At baseline, patients with documented penicillin allergy could be referred to the clinical pharmacist for evaluation by their family physician. To increase delabelling, a standardized letter was mailed to patients with documentation of penicillin allergy inviting them to schedule an appointment. At the appointment, the PEN-FAST tool was used to identify low-risk patients eligible for oral amoxicillin challenge. If the patient completed the oral amoxicillin challenge without anaphylaxis, the allergy was refuted on their chart. The project received exemption from the University of Saskatchewan Biomedical Research Ethics Board (E-Bio-091) and Operational Approval from the Saskatchewan Health Authority (OA-UofS-E-Bio-091).

Results/Findings: Five-hundred and eight FMU patients were identified as having a penicillin allergy. Of these patients, 102 (20%) were sent letters early February 2026 in a 2-month PDSA cycle. Initial phone visits booked were 15 (14.7%), patients delabelled were 10 (9.8%), 2 (13.3%) met exclusion criteria, and 3 (20%) have an upcoming challenge scheduled with results pending. Of patients who completed an oral amoxicillin challenge, 0 experienced an anaphylactic reaction. Since FMU began delabelling in December 2024, 5.4% of patients have been delabelled with an increase 1.9% during the first PDSA cycle.

Discussion: Overall, the project's aim was not met after one PDSA cycle with only 28 (5.4%) patients delabelled of those with documented allergy. Delabelling increased during the first PDSA cycle.

Conclusions: Additional strategies are required to encourage uptake from patients beyond a mailed letter.

Recommendations: Future PDSA cycles could focus on advising the patient’s family physician that a letter has been sent out to facilitate discussion recommending project uptake.

References:

1. Blumenthal KG, Peter JG, Trubiano JA, Phillips EJ. Antibiotic Allergy. *Lancet*. 2019 Jan;393(10167):183-198.
2. Salkind AR, Cuddy PG, Foxworth JW. Is This Patient Allergic to Penicillin: An Evidence-Based Analysis of the Likelihood of Penicillin Allergy. *JAMA*. 2001 May;285(19):2498-2505.
3. Powel N, Honeyforrd K, Sandoe J. Impact of Penicillin Allergy Records on Antibiotic Cost and Length of Hospital Stay: A Single-Centre Observational Retrospective Cohort. *J Hosp Infect*. 2020 Sep;106(1):35-42.
4. Copaescu AM, Vogrin S, James F, Chua KYL, Rose MT, De Luca J, et al. Efficacy of a Clinical Decision Rule to Enable Direct Oral Challenge in Patient with Low-Risk Penicillin Allergy: The PALACE Randomize Clinical Trial. *JAMA Intern Med*. 2023 Sep;183(9):944-952.
5. Cooper L, Harbour J, Sneddon J, Seaton RA. Safety and Efficacy of De-Labeling Penicillin Allergy in Adults Using Direct Oral Challenge: A Systematic Review. *JAC Antimicrob Resist* [Internet]. 2021 Jan 27 [cited 2026 May 14];3(1):[10 p.]. Available from: <https://doi.org/10.1093/jacamr/dlaa123>
6. Trubiano JA, Vogrin S, Chua KYL, Bourke J, Yun J, Douglas A, et al. Development and Validation of a Penicillin Allergy Clinical Decision Rule. *JAMA Intern Med* 2020 May;180(5):745-752.
7. Khan DA, Banerji A, Blumenthal KG, Phillips EJ, Solensky R, White AA, et al. Drug Allergy: A 2022 Practice Parameter Update. *J Allergy Clin Immunol*. 2022 Dec;150(6):1333-1393.
8. Saskatchewan Health Authority [Internet]. Regina (SK): Saskatchewan Health Authority; c2026. Firstline App; [cited 2026 May 14];[about 2 screens]. Available from: <https://library.saskhealthauthority.ca/home/mobile/firstline>.

Dance Movement Therapy's Role in Chronic Disease Management: A Systematic Review

Vladyslav Liamprekht, FMR II; Ai-Lien Le, FMR II; Vipul Parekh, MBBS, CCFP;
Rhonda Bryce, MD, MSc; Breanna Davis, MD, CCFP, FCFP

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Dance Movement Therapy (DMT) has been introduced to a variety of physical and mental health conditions in combination with standard models of care. With a growing interest in exercise prescribing across Canada, it is important to understand the benefits and challenges of DMT to better provide these programs to patients in primary care.

Question(s): What are the health outcomes of adding DMT in the management of chronic physical diseases compared to standard clinical care?

Methods/Methodology: This systematic review used the following databases: PubMed, Epistemonikos, and Cochrane Library. Only studies written in English, and which were published from 2000 onwards were included. Studies were excluded if the primary focus was a chronic mental health disease or had pediatric or cancer populations. For quality assessment, The Newcastle-Ottawa Scale, the Cochrane Risk of Bias Tool, and A Measurement Tool to Assess Systematic Reviews were used. Due to the situation of all study materials within the public domain, formal ethics approval was not required.

Results/Findings: Ten studies were identified which investigated physical and mental health outcomes and quality of life measures of DMT related to obesity, old age, diabetes, coronary artery disease, chronic heart failure, COPD and Parkinson's disease. The study design captured systematic reviews, randomized controlled trials, and prospective cohort studies. The included studies were conducted in a range of geographic regions: Canada, USA, Europe, Australia, Caribbean, and South Korea.

Discussion: Of the included studies, predominantly physical health outcomes were reported related to balance, gait, blood pressure, body weight and body fat – although it is unclear whether improved mobility translates to greater independence in daily activity or the lasting benefits of DMT. Improvements in self-esteem and body image were also noted compared to traditional activity training. In terms of feasibility, participants reported enjoyment and willingness to continue long term which is encouraging considering individualized exercise prescriptions.

Conclusions: The literature and other available data suggest that DMT has a beneficial role in chronic disease management with no reported adverse events. Further research is required to identify which DMT program would be most helpful for specific physical conditions.

References:

1. Duda-Sikuła M, Kurpas D. Enhancing Chronic Disease Management: Personalized Medicine Insights from Rural and Urban General Practitioner Practices. *J Pers Med* [Internet]. 2024 Jul 1 [cited 2026 May 12];14(7):[14 p.]. Available from: <https://doi.org/10.3390/jpm14070706>.
2. Lee IM, Paffenbarger RS Jr. Associations of Light, Moderate, and Vigorous Intensity Physical Activity with Longevity. The Harvard Alumni Health Study. *Am J Epidemiol*. 2000 Feb;151(3):293-9.
3. Fortier M, Morgan T, Tomasone J, Jain R. Integration of Exercise Professionals to Help Patients Adopt and Maintain Healthy Movement Behaviour: What Are We Waiting For? *Can Fam Physician*. 2024 Oct;70(10):611-613.
4. Wiedenhofer S, Koch SC. Active Factors in Dance/Movement Therapy: Specifying Health Effects of Non-Goal-Oriented Movement. *Arts Psychother*. 2017 Feb;52(2):10-23.
5. Liu W, Xue H, Wang Z. A Systematic Comparison of Intercultural and Indigenous Cultural Dance Education from a Global Perspective (2010-2024). *Front Psychol* [Internet]. 2024 Nov 26 [cited 2026 May 12];(15):[18 p.]. Available from: <https://doi.org/10.3389/fpsyg.2024.1493457>.
6. Zhang X, Wei Y. The Role of Dance Movement Therapy in Enhancing Emotional Regulation: A Literature Review. *Heliyon* [Internet]. 2024 Aug 3 [cited 2026 May 12];10(15):[17 p.]. Available from: <https://doi.org/10.1016/j.heliyon.2024.e35733>.
7. Santos D, Queiroz A, Menezes R, Bachion MM. Effectiveness of Senior Dance in the Health of Adults and Elderly People: An Integrative Literature Review. *Geriatr Nurs*. 2020 Sep-Oct;41(5):589-599.
8. Wshah A, Butler S, Patterson K, Goldstein R, Brooks D. “Let’s Boogie” Feasibility of a Dance Intervention in Patients with Chronic Obstructive Pulmonary Disease. *J Cardiopulm Rehabil Prev*. 2019 Sep;39(5):E14-E19.
9. Tomlinson CL, Patel S, Meek C, Herd CP, Clarke CE, Stowe R, et al. Physiotherapy Versus Placebo or No Intervention in Parkinson’s Disease. *Cochrane Databases Syst Rev* [Internet]. 2013 Sep 10 [cited 2026 May 12];2013(9):[110 p.]. Available from: <https://doi.org/10.1002/14651858.cd002817.pub4>.
10. Tomlinson CL, Herd CP, Clarke CE, Meek C, Patel S, Stowe R, et al. Physiotherapy for Parkinson’s Disease: A Comparison of Techniques. *Cochrane Databases Syst Rev* [Internet]. 2014 Jun 17 [cited 2026 May 12];2014(6):[108 p.]. Available from: <https://doi.org/10.1002/14651858.cd002815.pub2>.
11. Sherrington C, Fairhall NJ, Wallbank GK, Tiedemann A, Michaleff ZA, Howard K, et al. Exercise for Preventing Falls in Older People Living in the Community. *Cochrane Databases Syst Rev* [Internet]. 2019 Jan 31 [cited 2026 May 12];2019(1):[452 p.]. Available from: <https://doi.org/10.1002/14651858.cd012424.pub2>.
12. Negron C, Garcia MM, Sanchez G, Jovet-Toldeo GG, Martinez JA, Borrás IC. Effects of Tropical Latin Dance on Functional Capacity, Waist Circumference, and Quality of Life Compared to Traditional Aerobic Exercise Among Patients with Coronary Artery Disease in the VA Caribbean Healthcare System. *J Cardiopulm Rehabil Prev*. 2024 Sep;44(1):E68-E70.
13. Howe TE, Rochester L, Neil F, Skelton DA, Ballinger C. Exercise for Improving Balance in Older People. *Cochrane Databases Syst Rev* [Internet]. 2011 Nov 9 [cited 2026 May 12];2011(11):[267 p.]. Available from: <https://doi.org/10.1002/14651858.cd004963.pub3>.

14. Allet L, Muller-Pinget S, Punt I, Edelsten C, Ballif A, Golay A, et al. Dance Therapy Combined with Patient Education Improves Quality of Life of Persons with Obesity: A Pilot Feasibility Study for a Randomized Controlled Trial. *Obes Res Clin Pract*. 2017 Jan-Feb;11(1):79-87.
15. Ernst M, Folkerts AK, Gollan R, Lieker E, Caro-Valenzuela J, Adams A, et al. Physical Exercise for People with Parkinson's Disease: A Systematic Review and Network Meta-Analysis. *Cochrane Databases Syst Rev* [Internet]. 2024 Apr 8 [cited 2026 May 12];2024(4):[360 p.]. Available from: <https://doi.org/10.1002/14651858.cd013856.pub3>.
16. Kiepe MS, Stockigt B, Keil T. Effects of Dance Therapy and Ballroom Dance on Physical and Mental Illnesses: A Systematic Review. *Arts Psychother*. 2012 Nov;39(5):404-11.
17. Muller-Pinget S, Carrard I, Ybarra J, Golay A. Dance Therapy Improves Self-Body Image Among Obese Patients. *Patient Educ Couns*. 2012 Dec;89(3):525-28.

Enhanced Skills Program Evaluation: Graduate Survey of Career Satisfaction, Support Systems, and Work-Life Integration

Dinesh Kumar, FMR II; Maria Siddiqui, FMR II;
Martin Heroux, MD, CCFP, CCFP (EM), CCFP (SEM); Adam Clay, MSc

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Literature suggests that enhanced skills training can expand access to care but also highlights burnout, evolving practice patterns, and discrepancies between intended and actual scope of practice. Evaluation of the University of Saskatchewan's Enhanced Skills (USASK ES) programs ensures alignment with evolving community health needs and the delivery of comprehensive, patient-centered care.

Question(s): How do graduates of the USASK ES Programs apply their enhanced competencies in practice, and how are these patterns associated with job satisfaction and perceived support in their current work environments?

Methods/Methodology: An anonymous cross-sectional survey was conducted among graduates of USASK ES Programs. The survey included closed- and open-ended questions on key domains, including demographics, practice patterns, enhanced skill utilization, workplace environment, and job satisfaction. Quantitative data were analyzed using descriptive statistics, while open-ended responses were analyzed using thematic analysis. This project received approval from the University of Saskatchewan's Behavioural Research Ethics Board (Beh-REB #6022).

Results/Findings: Thirty-five responses were analyzed (response-rate 18%). Most respondents practiced in ES-related focused areas (85.7%), with 62.9% spending 76–100% of their clinical time in ES work. Only 34.3% reported practicing comprehensive family medicine, with marked differences between ESS/FPA (61.1%) and Emergency-Medicine (5.9%). Satisfaction was high, with 94.3% reporting that training met or exceeded expectations and 97% indicating they would pursue ES training again. Most (84.8%) reported improved ability to address community needs. Challenges included work-life balance (63.3%), difficulty balancing ES and generalist roles (46.7%), and call burden (36.7%). While 82.4% reported feeling supported, qualitative findings identified system-level challenges, including privileging barriers and variable specialist support.

Discussion: These findings align with literature showing ES training improves access to care but is associated with evolving practice patterns and wellness challenges. High satisfaction despite burnout risk aligns with prior studies and reflects focused practice, particularly in Emergency-Medicine. System-level barriers, including workload, call burden, and privileging challenges, influence ES use and highlight the need for improved mentorship, clearer integration, and better alignment of ES roles within healthcare systems.

Conclusions: ES training supports physician satisfaction and healthcare system capacity; however, variation in practice patterns and structural barriers impact integration into comprehensive family medicine and long-term sustainability.

References:

1. Grierson L, Vanstone M, Aliche I. Understanding the Impact of the CFPC Certificates of Added Competence [Internet]. Mississauga (ON): College of Family Physicians of Canada; 2020 Feb 6 [updated 2020 Mar 25]. 69 p. Available from: <https://www.cfpc.ca/CFPC/media/PDF/2020-04-CAC-Impact-Study-Report.pdf>.
2. Thommasen HV, Lavanchy M, Connelly I, Berkowitz J, Grzybowski S. Mental Health, Job Satisfaction, and Intention to Relocate. Opinions of Physicians in Rural British Columbia. *Can Fam Physician*. 2001 Apr;47(4):737-44.
3. Lee FJ, Stewart M, Brown JB. Stress, Burnout, and Strategies for Reducing Them: What's the Situation Among Canadian Family Physicians? *Can Fam Physician*. 2008 Feb;54(2):234-235.
4. Ringdahl E, Delzell JE Jr, Kruse RL. Changing Practice Patterns of Family Medicine Graduates: A Comparison of Alumni Surveys from 1998 to 2004. *J Am Board Fam Med*. 2006 Jul-Aug;19(4):404-412.
5. Coutinho AJ, Cochrane A, Stelter K, Phillips RL Jr, Peterson LE. Comparison of Intended Scope of Practice for Family Medicine Residents with Reported Scope of Practice Among Practicing Family Physicians. *JAMA*. 2015 Dec;314(22):2364-2372.
6. Canadian Medical Association. CMA 2021: National Physician Health Survey [Internet]. Ottawa (ON): Canadian Medical Association; 2022 Aug 24 [cited 2026 May 13]. 122 p. Available from: <https://digitallibrary.cma.ca/link/digitallibrary17>.
7. Canadian Institute for Health Information [Internet]. Ottawa (ON): Canadian Institute for Health Information; c2026. Access to primary care: many Canadians face challenges; 2024 Dec 5 [cited 2026 May 11];[about 9 screens]. Available from: <https://www.cihi.ca/en/primary-and-virtual-care-access-emergency-department-visits-for-primary-care-conditions/access-to-primary-care-many-canadians-face-challenges>.
8. Elma A, Vanstone M, Aliche I, Barber C, Howard M, Mountjoy M, et al. The Certificate of Added Competence Credentialing Program in Family Medicine: A Descriptive Survey of the Family Physician Perspective of Enhanced Skill Practices in Canada. *Can Med Educ J*. 2023 Nov 8;14(5):121-144.
9. College of Family Physicians of Canada. Residency Training Profile for Family Medicine and Enhanced Skills Programs Leading to Certificates of Added Competence [Internet]. Mississauga (ON): College of Family Physicians of Canada; 2021 May [cited 2026 May 11]. 77 p. Available from: <https://www.cfpc.ca/rtp>.

Pyxis-Programmed Anaphylaxis Kit to Improve Testing of Suspected Intraoperative Anaphylaxis: A Quality Improvement Project

Yechan Kim, FMR III (FPA); Vicky Loessin, MD, FRCPC;
Carla Flogan, RD, MSc; Richard Schaan, MD, FRCPC; Darcie Earle, BSc

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Intraoperative anaphylaxis is rare but potentially fatal. Confirming the diagnosis depends on timely serum tryptase sampling and proper documentation, yet these steps are often missed. National audit data show that guideline-concordant tryptase sampling is not consistently achieved. Without a standardized approach, diagnoses may be missed, triggers go unidentified, and patients face avoidable risk during future anesthetics.

Question(s): Does implementation of a Pyxis-programmed Anaphylaxis Kit improve adherence to recommended diagnostic testing, documentation, and follow-up planning for suspected intraoperative anaphylaxis? This project primarily addresses safety and effectiveness.

Methods/Methodology: This quality improvement project (PDSA Cycle 1) used a pre–post design in SHA operating rooms. The intervention was a Pyxis-programmed Anaphylaxis Kit bundling key emergency medications (epinephrine, diphenhydramine, famotidine, methylprednisolone) with an embedded cognitive aid checklist. Cases were identified through anesthesiologist self-report and laboratory records. Data collected included tryptase ordering, timing, documentation, and follow-up. An anonymous survey of pediatric anesthesiologists (10/16, 62.5%) assessed awareness, usefulness, and suggestions for improvement. This project was granted exemption by the University of Saskatchewan Research Ethics Board (E-BIO-111).

Results/Findings: Pre-implementation review identified 3 cases: tryptase was ordered and documented in 2/3 (67%). One case had no tryptase ordered, no anaphylaxis documented, and no follow-up plan. One post-implementation case has occurred so far—tryptase was ordered and anaphylaxis was documented, though some data fields are still pending. On the survey, 100% rated the kit as useful (50% extremely, 40% very, 10% somewhat), 100% agreed the medications were appropriate, and 90% wanted a weight-based dosing chart included.

Discussion: Because intraoperative anaphylaxis is uncommon (~5 cases/year locally), early evaluation focuses on usability and adoption rather than clinical outcomes. Survey responses suggest the kit helps standardize crisis management and reduces cognitive burden. The inconsistency seen in pre-implementation data supports the rationale for this kind of structured approach.

Conclusions: A Pyxis-programmed Anaphylaxis Kit is a practical, well-received intervention that can support the management of intraoperative anaphylaxis at the point of care.

Recommendations: Next PDSA cycles should add a weight-based dosing chart and tryptase blood tubes based on survey feedback, with continued data collection and expansion to other SHA sites.

Investigating Hypocalcemia in Trauma Patients: Trends in Detection and Treatment Timelines at RUH

Chiranjeevi Gade, FMR III; Troy Simpson, FMR III; Matt Johnson, MD, FPA, FRCPC

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Hypocalcemia is common in critically ill trauma patients and contributes to coagulopathy, impaired cardiac contractility, and poor outcomes. It may result from hemorrhage, citrate chelation during transfusion, and acid–base disturbances. Despite its clinical importance, the timing of detection and adequacy of calcium replacement in acute trauma care remain unclear.

Question(s): What is the local incidence of hypocalcemia in Level 1 trauma patients at Royal University Hospital (RUH)? What are the timelines for its identification and treatment? How is the effectiveness of calcium replacement to improve efficiency evaluated?

Methods/Methodology: A retrospective chart review of 150 Level 1 trauma patients presenting to RUH between July 2024 and August 2025 was conducted. Data were extracted from electronic and paper records, including laboratory timelines, transfusion data, and medication administration records. Injury severity was categorized using the Injury Severity Score (ISS). Primary outcomes included incidence of hypocalcemia, time to detection, time to calcium administration, and persistence of hypocalcemia after treatment. Approval from University of Saskatchewan’s REB Bio #5946 and SHA OA-UofS-5946.

Results/Findings: Hypocalcemia was identified in 26/150 patients (18%) on presentation. Among these, 20 patients received calcium replacement; however, all remained hypocalcemic afterward. Fourteen patients with hypocalcemia received no treatment. Notably, hypocalcemia preceded transfusion in 11 cases. Laboratory turnaround times varied significantly (lab order to reporting: 21–120 minutes total). Blood transfusion was more common in major trauma (ISS >15), with 31/65 such patients receiving blood. Calcium dosing ranged from 1–9 g of calcium gluconate. One patient presented with hypercalcemia and died.

Discussion: Hypocalcemia occurs early in trauma and is not solely transfusion-related. Delays in laboratory reporting and inconsistent repeat testing may contribute to under-recognition. Current replacement strategies appear inadequate, as most treated patients remained hypocalcemic. These findings highlight a gap in timely correction and suggest that earlier, protocolized calcium administration may be necessary.

Conclusions: Hypocalcemia is frequent and often undertreated in trauma patients at RUH. Improvements in rapid detection and standardized calcium replacement—potentially integrated into massive transfusion protocols—may enhance patient outcomes.

Do Visual Reminders Improve Appropriate ASA Prescribing for Preeclampsia Prevention?: A Quality Improvement Study

Yuli Chen, FMR II; Calista Lytle, FMR II; Audrey Roane, FMR II;
Breanna Davis, MD, CCFP, FCFP; Rhonda Bryce, MSc, MD;
Vivian R Ramsden RN, PhD, MCFP (Hon.)

Department of Family Medicine, College of Medicine, University of Saskatchewan

ABSTRACT

Background: Current guidelines by the Society of Obstetricians and Gynecologists of Canada indicate low-dose ASA for preeclampsia prevention for patients with a BMI ≥ 30 (2). In Northern Saskatchewan, obesity rates are among the highest in Canada (1), and providers have noted many patients presented beyond the window to initiate ASA. This project aims to reduce the negative impact of preeclampsia by improving early ASA initiation.

Question(s): In pregnant patients with a pre-pregnancy or early pregnancy BMI ≥ 30 receiving early prenatal care, does the implementation of a prenatal form visual reminder increase the proportion of patients appropriately prescribed ASA?

Methods/Methodology: A visual prompt was added to the prenatal record to encourage screening for BMI to initiate ASA. Chart reviews compared patients four months pre-intervention and four months post-intervention. As a continuation of a previous quality improvement research project, this project was exempted from the requirements of Research Ethics Board (REB), E-Bio-075.

Results/Findings: Among 139 patients (106 pre-intervention, 33 post-intervention), 39% had an ASA indication. Overall, 85% of patients presented before 16 weeks. Out of these, 18% did not have a prenatal record prior to 16 weeks. In addition, 18% of the total cohort lacked baseline BMI documentation representing a primary barrier to care. Among patients with an ASA indication presenting before 16 weeks, appropriate prescribing declined from 72.0% pre-intervention to 33.3% post-intervention ($p=0.23$). Patients with an ASA indication who received ASA had lower preeclampsia rates (8.7%) compared to untreated patients (33.3%; $p=0.15$).

Discussion: While most patients presented within the treatment window prior to 16 weeks, BMI was frequently undocumented and the prenatal record was often not utilized at those visits. Furthermore, the low sample size in the post-intervention group limits statistical power to determine benefit.

Conclusions: Passive visual reminders did not demonstrate a benefit in increasing appropriate ASA prescriptions for the prevention of preeclampsia.

Recommendations:

1. Request medical office assistants to capture height and weight during positive urine pregnancy tests or initial prenatal workups.

2. Keep the visual prompt and continue encouraging providers to use the prenatal record by 12 weeks gestation.

References:

1. Dzakpasu S, Nelson C, Darling EK, Edwards W, Murphy PA, Scott H, et al. Trends in Rate of Hypertensive Disorders of Pregnancy and Associated Morbidities in Canada: A Population-Based Study (2012-2021). *CMAJ*. 2024 Jul;196(26):E897–904.
2. Lim S, Li W, Kemper J, Nguyen A, Mol BW, Reddy M. Biomarkers and the Prediction of Adverse Outcomes in Preeclampsia: A Systematic Review and Meta-analysis. *Obstet Gynecol*. 2021 Jan;137(1):72–81.
3. Cadavid AP. Aspirin: The Mechanism of Action Revisited in the Context of Pregnancy Complications. *Front Immunol* [Internet]. 2017 Mar 15 [cited 2026 May 15];8:[8 p.]. Available from: <https://doi.org/10.3389/fimmu.2017.00261>.
4. Chaddha V, Viero S, Huppertz B, Kingdom J. Developmental Biology of the Placenta and the Origins of Placental Insufficiency. *Semin Fetal Neonatal Med*. 2004 Oct;9(5):357–369.
5. Roberge S, Bujold E, Nicolaides KH. Aspirin for the Prevention of Preterm and Term Preeclampsia: Systematic Review and Meta-Analysis. *Am J Obstet Gynecol*. 2018 Mar;218(3):287-293.
6. Roberts JM. Pathophysiology of Ischemic Placental Disease. *Semin Perinatol*. 2014 Apr;38(3):139–145.
7. Rolnik DL, Nicolaides KH, Poon LC. Prevention of Preeclampsia with Aspirin. *Am J Obstet Gynecol*. 2022 Feb;226(2S):S1108–1119.
8. Rolnik DL, Wright D, Poon LC, O’Gorman N, Syngelaki A, de Paco Matallana C, et al. Aspirin Versus Placebo in Pregnancies at High Risk for Preterm Preeclampsia. *N Engl J Med*. 2017 Aug 17;377(7):613–622.
9. Garza-Galvan ME, Ferrigno AS, Campos-Zamora M, Bain PA, Easter SR, Kim J, et al. Low-Dose Aspirin Use in the First Trimester of Pregnancy and Odds of Congenital Anomalies: A Meta-Analysis of Randomized Controlled Trials. *Int J Gynaecol Obstet*. 2023 Feb;160(2):526–537.
10. Magee LA, Smith GN, Bloch C, Côté AM, Jain V, Nerenberg K, et al. Guideline No. 426: Hypertensive Disorders of Pregnancy: Diagnosis, Prediction, Prevention, and Management. *J Obstet Gynaecol Can*. 2022 May;44(5):547-571.
11. Magee LA, Pels A, Helewa M, Rey E, Von Dadelszen P. Diagnosis, Evaluation, and Management of the Hypertensive Disorders of Pregnancy. *Pregnancy Hypertens*. 2014 Apr;4(2):105–145.
12. Horgan R, Hage Diab Y, Waller J, Abuhamad A, Saade G. Low-Dose Aspirin Therapy for the Prevention of Preeclampsia: Time to Reconsider Our Recommendations? *Am J Obstet Gynecol*. 2023 Oct;229(4):410–418.

13. Maxwell C, Gaudet L, Cassir G, Nowik C, McLeod NL, Jacob CÉ, et al. Guideline No. 391-Pregnancy and Maternal Obesity Part 1: Pre-Conception and Prenatal Care. *J Obstet Gynaecol Can.* 2019 Nov 1;41(11):1623–40.
14. Public Health Agency of Canada. Obesity Statistics in Canada [Internet]. Ottawa (ON): Government of Canada; 2024 Sep [cited 2026 May 14]. 120 p. Available from: [https://www.canada.ca/content/dam/phac-aspc/documents/services/publications/ healthy-living/obesity-statistics-canada/obesity-statistics-canada.pdf](https://www.canada.ca/content/dam/phac-aspc/documents/services/publications/healthy-living/obesity-statistics-canada/obesity-statistics-canada.pdf).
15. Salmon C, Bordun-Slater KA. Increasing Acetylsalicylic Acid Prescribing in Pregnancy Based on the New SOGC Guidelines: A Program Evaluation in Prince Albert, Saskatchewan. Paper presented at: Department of Academic Family Medicine’s 34th Annual Resident Scholarship Day; 2024 May 31; Saskatoon, SK.
16. Arditi C, Rège-Walther M, Durieux P, Burnand B. Computer-Generated Reminders Delivered on Paper to Healthcare Professionals: Effects on Professional Practice and Healthcare Outcomes. *Cochrane Database Syst Rev* [Internet]. 2017 Jul 6 [cited 2026 May 14];7(7):[91 p.]. Available from: <https://doi.org/10.1002/14651858.cd001175.pub4>.
17. American College of Obstetricians and Gynecologists. Committee Opinion No. 680 Summary: The Use and Development of Checklists in Obstetrics and Gynecology. *Obstet Gynecol* 2016 Nov;128(5):1200.
18. Statistics Canada [Internet]. Ottawa (ON): Government of Canada; c2026. Profile table, Census Profile, 2021 Census of Population - Prince Albert, City (CY) [Census subdivision], Saskatchewan [Internet]. 2022 [cited 2026 May 10];[about 68 screens]. Available from: <https://www12.statcan.gc.ca/census-recensement/2021/dp-pd/prof/details/page.cfm?Lang=E&SearchText=prince%20albert&DGUIDlist=2021A00054715066&GENDERlist=1,2,3&STATISTIClist=1,4&HEADERlist=0>.
19. Health Canada. Prenatal Nutrition Guidelines for Health Professionals: Gestational Weight Gain [Internet]. Ottawa (ON): Government of Canada; 2010 [cited 2026 May 12]. 21 p. Available from: https://www.canada.ca/content/dam/hc-sc/migration/hc-sc/fn-an/alt_formats/pdf/nutrition/prenatal/ewba-mbsa-eng.pdf.